

- AAC. *See* Aortic annular calcification
- Abciximab
 adjunctive, and PCI, ST segment analysis in AMI after, 48
 effect on myocardial reperfusion after PCI for STEMI, 798
 effects on platelet inhibition, 798
 in patients undergoing high-risk PCI, 489
 for thrombocytopenia, 426
- Ablation. *See also* Catheter ablation; Septal ablation
 of His bundle, left ventricular remodeling during long-term right ventricular pacing after, 1223
 pulmonary vein, use of esophageal leads in determining strategies of, 1494
 slow pathway, effects on electrophysiologic characteristics of atrioventricular nodal reentry tachycardia, 860
- ACC/AHA. *See* American College of Cardiology/American Heart Association
- ACE inhibitors. *See* Angiotensin-converting enzyme inhibitors
- ACS. *See* Acute coronary syndrome
- Activated clotting time, optimal, during carotid artery stenting, 1657
- Acute coronary syndrome
 drawbacks of nonrandomized clinical trials, 151 (Readers' Comments)
 intravascular ultrasound of coronary arteries during PCI in, 29
 non-ST-elevation
 effects of drug-eluting vs bare metal stents on biomarkers after PCI for, 1473
 frequency and consequences of electrocardiogram recording >10 minutes after ER arrival, 437
 impact of CHF, 1707
 prediction of long-term mortality by biomarkers, 167
 troponin T and ST segment in lead aVR in, 334
 relation of blood glucose to long-term mortality in, 1573
 serum levels of inflammatory markers in patients with
 and coronary vasospasm without CAD, 1429
 and stable angina with CAD, 1429
- Acute myocardial infarction
 after PCI
 and adjunctive abciximab, ST segment analysis, 48
 effect of reactive hyperemia on myocardial tissue reperfusion in, 617
 left ventricular systolic vs diastolic function as outcome predictor, 160
 prognostic effects of distal vs proximal coronary occlusion, 1137
 angiographic no-reflow phenomenon during stent implantation, 301
 anterior wall, and after PCI, coronary microvascular resistance during, 1131
 in apparently healthy subjects aged ≥ 55 years, relation to ventricular arrhythmias, 1351
 coronary thrombus of patients with
 fibrin in, 26
 platelets in, 26
 tissue factor in, 26
 von Willebrand factor in, 26
 effect of β -blockers on survival after, 1125 (Readers' Comments)
 effects of optimized medical and revascularization therapy on incidence of sudden cardiac death after, 480
 effects of statins on CHF and myocardial ischemia in Japanese patients with, 1165
 gender differences of revascularization in, 1722
 mortality prediction using
 QRS wave, 353
 T wave, 353
 N-terminal ProBNP on patient admission, relation to scintigraphic infarct size, reperfusion efficacy and prognosis, 1151
 outcomes
 after balloon angioplasty vs stent implantation, 977
 in-hospital, prediction by white blood cell count and plasma glucose, 1558
 prediction by aortic root dimension in persons ≥ 65 years of age, 270
 of President Eisenhower in 1955, 1125 (Readers' Comments)
 relation of B-type natriuretic peptide to left ventricular diastolic function and extent of myocardial scar early after, 1146
 return to full normal activities including work at 2 weeks after, 952
 risk of recurrence, relation to statin initiation after hospital discharge in patients ≥ 65 years of age, 115
 routine admission laboratory tests to predict thirty-day mortality in, 1435
 SPECT vs myocardial contrast echocardiography to predict transmural of, 1718
 vs unstable angina pectoris, vs stable angina pectoris, serum MMP-9 in, 175
- Acute pulmonary embolism
 bedside testing for brain natriuretic peptides to identify right ventricular dysfunction and outcome in, 1386
 predictors of hemorrhage after fibrinolysis for, 127
- Adiponectin, in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646
- Adolescents, incidence and risk factors of heart failure in congenital heart disease after cardiac surgery, 1238
- Adults
 with CAD, and with vs without metabolic syndrome, effect of simvastatin on outcomes of, 1487
 incidence and risk factors of heart failure in congenital heart disease after cardiac surgery, 1238
 with operated tetralogy of Fallot, relation of biventricular dysfunction to elevated NT-pro-BNP and reduced exercise capacity, 1377
- Aerobic exercise. *See* Exercise
- African-Americans
 admitted for CHF at a university teaching hospital, quality of care in, 690
 vs Caucasians, phenotypic predictors of response to simvastatin therapy, 843
 efficacy and safety of rosuvastatin vs atorvastatin for hypercholesterolemia, 229
 men and women, blood pressure stress reactivity and left ventricular mass among, 240
 vs non-African-Americans, with chronic heart failure, metabolic vasodilation in response to exercise, ischemia, and endothelium-dependent flow-mediated dilation, 685
- Aged
 50 to 79 years, QT interval and subintervals from ethnically diverse population of women, 730
 70 to 79 years, relation of social integration to CRP and IL-6 levels in men and women, 1010
 ≥ 65 years
 aortic root dimension in predicting heart failure, stroke, cardiovascular mortality, all-cause mortality and AMI, 270
 left atrium in systolic vs diastolic heart failure, 83
 time of statin initiation after hospital discharge, relation to mortality and risk of recurrence in AMI patients, 155
 >65 years, relation of ACE inhibitor treatment to IGF-I serum levels, 1525
- Air pollution, relation to hospital admissions for CHF in seven United States cities, 404
- Alcohol
 septal ablation, for obstructive hypertrophic cardiomyopathy
 predictors of subacute complete heart block after, 264
 time course of pressure gradient response after, 1511
- Aldosterone
 levels in coronary sinus and ascending aorta
 of aortic stenosis patients, 1068, 1107 (Editorial)
 of CHD patients, 1068, 1107 (Editorial)
 plasma levels, for prediction of in-stent restenosis, 785
- American College of Cardiology/American Heart Association, guidelines on periprocedural anticoagulation in patients with mechanical valves, adherence to, 891
- American Journal of Cardiology*, proceedings of the Editorial Board meeting on 13 March 2006, 1670 (From the Editor)
- AMI. *See* Acute myocardial infarction
- Amiodarone, causing severe ataxia, 1550 (Readers' Comments)
- Amplatzer septal occluder, cardiac conduction disturbances and ventricular arrhythmias after transcatheter closure of muscular ventricular septal defects, 558

- Androgenic anabolic steroids, effects on left ventricular diastolic function in male body builders, 912
- Aneurysm
- abdominal aortic
 - and cardiovascular disease in men, 1123 (Readers' Comments)
 - risk in CAD, peripheral arterial disease as marker of, 1549 (Readers' Comments)
 - size of, and mortality in patients treated with and without statins, 279
 - atrial septal, familial origin of, 148
 - left ventricular apical, and hypertrophic cardiomyopathy in identical twins, 1109
- Angina pectoris
- pre-MI, effect on reperfusion arrhythmias, 1157
 - refractory
 - transendocardial autologous bone marrow transplantation for, 823
 - two-year outcomes after EECF, 17
 - severe, intramyocardial injection of autologous bone marrow-derived mononuclear cells for, 1326
 - stable
 - and ACS with CAD, vs coronary vasospasm without CAD, serum levels of inflammatory markers in patients with, 1429
 - angiogenic factors in, 472
 - effects of high vs low clopidogrel loading dose on patients subjected to PCI, 984
 - soluble CD40 ligand in prediction of stent restenosis, 198
 - and unstable, safety and efficacy of cobalt-chromium alloy-based stent for, 349
 - vs unstable, vs AMI, serum MMP-9 in, 175
 - unstable
 - effects of morning vs evening atorvastatin intake after first elective PCI, 44
 - impact of clopidogrel on circulating levels of soluble CD40 ligand in patients undergoing coronary stenting, 192
 - relation to heart rate variability to serum CRP, 1702
 - vasospastic, intravascular ultrasound of coronary vasospasm induced by ergonovine maleate in, 1322
- Angiogenesis, therapeutic, in CHD, placebo effect on, 152 (Readers' Comments)
- Angiogenic factors
- relation to development of collateral vessels, angiographic evaluation, 472
 - in stable angina pectoris and total coronary artery occlusion, 472
- Angiography. *See also* Coronary angiography
- adverse events during PCI, relation to CK-MB elevation and ischemic end points, 1591
- AMI
- after PCI, prognostic effects of distal vs proximal coronary occlusion on, 1137
 - no-reflow phenomenon during stent implantation in, 301
- CAD
- contrast-enhanced MRI in detection of myocardial scar in, 768
 - gender differences in relation of renal impairment to long-term mortality and, 630
 - relation of ascending aortic pressures to outcomes in patients with, 590 (Readers' Comments)
 - relation of carotid intima-media thickness to severity and extent of, 624
 - severity in diabetes mellitus, relation to hemoglobin A_{1c} levels, 968
 - early outcome of ostial de novo left anterior descending coronary artery lesions treated with drug-eluting stents, 187
 - multidetector row spiral computed tomography for detection of significant coronary artery stenosis, 343
 - relation of development of collateral vessels to angiogenic factors, 472
 - restenosis after implantation of overlapping sirolimus-eluting (Cypher) stents, 499
 - safety and efficacy of cobalt-chromium alloy-based stent for stable and unstable angina pectoris, 349
 - transfemoral coronary, manual pressure hemostasis after, effectiveness in patients on warfarin anticoagulation, 485
- Angioplasty. *See also* Percutaneous transluminal coronary angioplasty balloon
- vs stent implantation, outcomes of AMI after, 977
 - vs vascular brachytherapy vs drug-eluting stents, safety and efficacy for saphenous vein graft in-stent restenosis, 1303
- primary coronary, vs fibrinolysis, ST-segment elevation for predicting outcomes after, 611
- Angioscopy, serial, after intracoronary brachytherapy for in-stent restenosis, 21
- Angiotensin-converting enzyme inhibitors
- effect on infarct-related coronary artery patency, 7
 - relation to IGF-I serum levels in subjects >65 years of age, 1525
 - use before STEMI, 7
- Angiotensin II, levels in coronary sinus and ascending aorta
- of aortic stenosis patients, 1068, 1107 (Editorial)
 - of CHD patients, 1068, 1107 (Editorial)
- Angiotensin receptor blockers, in prevention and treatment of arrhythmias, 921
- Ankle-brachial index, relation to coronary calcium, 181
- Anticoagulation
- over, warfarin-related, in patients with mechanical heart valves
 - FFP for, 409
 - vitamin K for, 409
 - periprocedural, in patients with mechanical valves, adherence to ACC/AHA guidelines on, 891
 - warfarin, effectiveness of manual pressure hemostasis after transfemoral coronary angiography in patients on, 485
- Aorta
- ascending
 - of aortic stenosis and CHD patients, levels of aldosterone, angiotensin II and B-type natriuretic peptides in, 1068, 1107 (Editorial)
 - pressures of, relation to outcomes in patients with angiographically demonstrated CAD, 590 (Readers' Comments)
 - distensibility and dimensions, in Turner syndrome, effects of growth hormone treatment, 1644
 - elastic properties of, in children after tetralogy of Fallot repair, 905
 - wall mechanics, in Marfan syndrome, assessment by transesophageal tissue Doppler echocardiography, 571
 - Aortic aneurysm, abdominal
 - and cardiovascular disease in men, 1123 (Readers' Comments)
 - risk in CAD, peripheral arterial disease as marker of, 1549 (Readers' Comments)
 - size of, and mortality in patients treated with and without statins, 279
 - Aortic annular calcification, relation to morbidity and mortality in community-dwelling elderly individuals, 1281
 - Aortic cross clamping, during major vascular surgery, effects of statins on renal function after, 1383
 - Aortic regurgitation, in children after tetralogy of Fallot repair, 905
 - Aortic root
 - dilation, in children after tetralogy of Fallot repair, 905
 - dimension in persons \geq 65 years of age, in prediction of all-cause and cardiovascular mortality, 270
 - AMI, 270
 - heart failure, 270
 - stroke, 270
 - distensibility vs diameter, as risk factor in Marfan syndrome, 1422 (Readers' Comments)
 - Aortic valve
 - calcification, relation of inflammation to risk factors of, 1502
 - heaviest known operatively-excised, 588
 - Aortic valve sclerosis, relation to morbidity and mortality in community-dwelling elderly individuals, 1281
 - Aortic valve stenosis
 - coronary sinus and ascending aorta levels of aldosterone, angiotensin II and B-type natriuretic peptide, 1068, 1107 (Editorial)
 - CRP as predictor of progression of, 90
 - apo-A-I. *See* Apolipoprotein A-I
 - apo-B. *See* Apolipoprotein B
 - Apolipoprotein A-I, effect on atherogenic particle number in predicting risk of fatal MI, 943
 - Apolipoprotein B
 - effect on atherogenic particle number in predicting risk of fatal MI, 943
 - and long-term risk of CHD in men, 997
 - phenotypic predictors of response to simvastatin therapy among African-Americans vs Caucasians, 843
- Arrhythmias

- after reperfusion, effect of pre-MI angina pectoris, 1157
 caused by implantable cardioverter-defibrillators, 1255
 prevention and treatment with angiotensin receptor blockers, 921
 recurrence during pregnancy in women with previous tachyarrhythmia, impact on fetal and neonatal outcomes, 1206
- ventricular
 after transcatheter closure of muscular ventricular septal defects using Amplatzer occluder, 558
 in CAD, effect of statins on risk of, 153 (Readers' Comments)
 high risk in patients with nonischemic dilated cardiomyopathy with syncope, 416
 relation to all-cause mortality and AMI in apparently healthy subjects aged ≥ 55 years, 1351
 worsening, in idiopathic dilated cardiomyopathy, prognostic value of CRT criteria for, 393
- Arrhythmogenic right ventricular dysplasia
 in children, MRI diagnosis of, 1232
 feasibility and variability of three-dimensional echocardiography vs MRI in, 703
- Arterial stiffness, effects of quinapril, 916
- Ascending aorta. *See* Aorta
- Asian Indians
 prediction of low-density lipoprotein particle size, phenotype B and particle concentrations by triglyceride/high-density lipoprotein ratio, 1007
 relation of lipoprotein lipase polymorphism to dyslipidemia and diabetes, 1337
- Aspirin
 and clopidogrel
 effects on platelet function, 38
 long-term use after PCI, 38
 effect on infarct-related coronary artery patency, 7
 influence on length of hospitalization in major vascular surgery patients, 1423
 reply to a reader's comment, 294
 resistance to, in type 1 vs type 2 diabetes, 567
 sensitivity, rapid desensitization for parents with, 294 (Readers' Comments)
 use before STEMI, 7
- Ataxia, severe, caused by amiodarone, 1550 (Readers' Comments)
- Atherosclerosis
 coronary
 characterization using intravascular ultrasound vs multidetector computed tomography, 598
 relation to left ventricular hypertrophy, effect of ethnicity, 1441
 vs saphenous vein grafts, intravascular ultrasound of, 593
 development of, influence of lipids, 372
 particle size, in predicting risk of fatal MI
 effects of apo-A-I, 943
 effects of apo-B, 943
 effects of low-density lipoprotein, 943
- Atherosclerotic vascular disease. *See* Coronary artery disease; Peripheral arterial disease; Stroke
- Athletes, male bodybuilders, effects of androgenic anabolic steroids on left ventricular diastolic function in, 912
- Atorvastatin
 benefits in CAD, effects of obesity, 1553
 combination with rosiglitazone, vs rosiglitazone monotherapy, effects on biomarkers of cardiovascular disease in type 2 diabetes, 646
 comparative safety of 80 mg vs 10 mg, 61
 effect on recurrence of atrial fibrillation after electrical cardioversion, 1490
 effects on endothelial function in hyperlipidemia, 1002
 for heart failure secondary to inflammatory dilated cardiomyopathy and elevated low-density lipoprotein cholesterol levels, 899
 morning vs evening intake, effect on major cardiac events and restenosis rates after first elective PCI, 44
 vs rosuvastatin, efficacy and safety in hypercholesterolemic African-Americans, 229
- Atrial appendage, left, thrombus detection by transesophageal echocardiography 421
- Atrial fibrillation
 after CABG, effect of statins on risk of, 55, 1266 (Readers' Comments), 1551 (Readers' Comments)
 frequency in mitral regurgitation secondary to mitral valve prolapse, 1039
 incidence of, effects of CRP, complement C3 and C4, 245
 incidence of cerebral ischemia in, role of hyperviscosity and inflammation, 1745
 before isolated mitral valve replacement for rheumatic mitral stenosis, 1035
 left atrial reverse remodeling after catheter ablation for, 1220
 lone, first onset and recurrence, relation to high-sensitivity CRP, 659
 in mitral stenosis after PBMV, one-year outcome of cardioversion of, 1045
 new-onset, patterns and predictors of warfarin use in patients with, 538
 with or without hypertension, CRP in, 1346
 paroxysmal, paroxetine for depressed men with, 1749
 persistent, before and after cardioversion, atrial and ventricular volume and function in, 1213
 plasma atrial natriuretic peptide and brain natriuretic peptide levels after catheter ablation of, 1741
 recurrence after electrical cardioversion, effect of atorvastatin, 1490
 selective inhibition of MMP and TIMPs in atrial and ventricular myocardium in patients with, 532
 time to first recurrence, relation to burden, 1343
- Atrial natriuretic peptide. *See* Natriuretic peptides
- Atrial remodeling, left
 effects of quinapril, 916
 reverse, after catheter ablation for atrial fibrillation, 1220
- Atrial septal aneurysm, familial origin of, 148
- Atrioventricular annulus, proximity to coronary arteries in young patients, implications in catheter ablation, 1752
- Atrioventricular delay
 in CRT, Doppler optimization of, relation to invasive hemodynamics in heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, 552
 echocardiographic evaluation, with dual-chamber pacing, long-term effects on hypertrophic cardiomyopathy, 1769
- Atrioventricular nodal reentry tachycardia
 effects of slow pathway ablation on electrophysiologic characteristics of, 860
 gender differences in electrophysiologic parameters, 384
- Atrium. *See also* Left atrium; Right atrium
 decreased function, contribution in pathogenesis of neurally mediated syncope, 1017
 myocardium in patients with atrial fibrillation, selective induction of TIMPs and MMP in, 532
 right-to-left shunting, diagnosis with transesophageal vs transthoracic echocardiography, 434 (Readers' Comments)
 volume and function, in persistent atrial fibrillation before and after cardioversion, 1213
- Autologous bone marrow-derived mononuclear cells, intramyocardial injection
 for severe angina pectoris, 1326
 for stress-induced myocardial ischemia, 1326
- Autologous bone marrow transplantation, transcatheter
 for myocardial ischemia, 823
 for refractory angina, 823
- AVS. *See* Aortic valve sclerosis
- Balloon dilatation
 of peripheral pulmonary stenosis, efficacy of medium-sized cutting balloons vs standard balloons, 1060
 predilatation, effectiveness of "direct" stenting without, 316
- Balloon valvuloplasty, percutaneous mitral, one-year outcome of cardioversion of atrial fibrillation in mitral valve stenosis after, 1045
- Barnard, Christiaan, 435 (Readers' Comments)
- β -blockers. *See* Beta-adrenergic blocking agents
- Behçet's disease, pulse-wave Doppler and Doppler tissue imaging of left ventricular function in, 716
- Beta-adrenergic blocking agents
 dobutamine stress echocardiography in observing wall motion abnormalities during recovery after administration of, 462
 effect on infarct-related coronary artery patency, 7
 effect on survival after AMI, 1125 (Readers' Comments)
 influence on length of hospitalization in major vascular surgery patients, 1423
 use before STEMI, 7
- Bifurcation lesions, coronary, long-term outcomes of sirolimus-eluting stent implantation by crush technique, 1317
- Biomarkers

- after PCI for non–ST-elevation ACS, effects of drug-eluting vs bare metal stents, 1473
- of cardiovascular disease, in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646
- lipid and lipoprotein, for CAD, 640
- for predicting long-term mortality
- in diabetes mellitus, 167
 - in non–ST-elevation ACS, 167
- Blalock, Alfred, 435 (Readers' Comments)
- Blood flow, cerebral, decreased in CHF secondary to idiopathic dilated cardiomyopathy, 1365
- Blood flow velocity, diastolic septal, in hypertrophic cardiomyopathy vs hypertensive left ventricular hypertrophy, 106
- Blood pressure
- ascending aortic, relation to outcomes in patients with angiographically demonstrated CAD, 590 (Readers' Comments)
 - effects of intravenous vs intrarenal fenoldopam in patients undergoing cardiac catheterization, 1076
 - in healthy postmenopausal women, effects of vegetarian diet, 380
 - reactivity to stress, and left ventricular mass among Caucasian and Africa-American men and women, 240
 - rising, effects on cardiovascular disease in China, 1676 (Readers' Comments)
- Body mass index
- effect on left ventricular cavity size and ejection fraction, 725
 - impact on systemic hypertension in lean Chinese, 839
 - increased
 - among Hispanic women, 68
 - and impaired endothelial function, 68
 - and leukocyte count, relation to metabolic syndrome, 835
 - relation to CAD severity in patients undergoing coronary angiography, 1277
- Bone marrow–derived mononuclear cells, autologous, intramyocardial injection for severe angina pectoris, 1326
- for stress-induced myocardial ischemia, 1326
- Bone marrow transplantation, autologous transendocardial
- for myocardial ischemia, 823
 - for refractory angina, 823
- Brachial artery, flow-mediated dilation
- circadian variation in idiopathic dilated cardiomyopathy, 699
 - in hyperlipidemia, effects of atorvastatin, 1002
- Brachytherapy
- vs drug-eluting stent implantation, intravascular ultrasound of restenosis after, 1292
 - intracoronary, for in-stent restenosis, serial angiography after, 21
 - vascular, vs drug-eluting stents, vs balloon angioplasty, safety and efficacy for saphenous vein graft in-stent restenosis, 1303
- Brain natriuretic peptides. *See* Natriuretic peptides
- Breath tests, carbon monoxide to measure influence of prosthetic heart valves on erythrocyte survival, 1374
- Brewer, Hollis Bryan, Jr., 1791 (Conversation With the Editor)
- Bundle branch block, left, with or without type 2 diabetes, extent of CAD in, 857
- Bundle of His, left ventricular remodeling during long-term right ventricular pacing after ablation of, 1223
- CABG. *See* Coronary artery bypass grafting
- CAD. *See* Coronary artery disease
- Calcification, aortic and mitral valves
- relation to morbidity and mortality in community-dwelling elderly individuals, 1281
 - risk factors of, relation to inflammation, 1502
- Calcium, coronary
- relation to ankle-brachial index, 181
 - relation to CHD, 1564
 - relation to left ventricular hypertrophy, effect of ethnicity, 1441
 - vs stress myocardial perfusion imaging, in apparently healthy siblings of individuals with premature CAD, 328
- Cancer, transesophageal echocardiography in detection of
- cardiac sources of embolism in, 894
 - nonbacterial thrombotic endocarditis in, 894
- Carbon monoxide, breath, to measure influence of prosthetic heart valves on erythrocyte survival, 1374
- Cardiac catheterization
- effects of intravenous vs intrarenal fenoldopam on kidney function and blood pressure in patients undergoing, 1076
 - laboratory, and patient transfer for PCI, 591 (Readers' Comments)
- Cardiac output, gain from CRT in patients with CAD or idiopathic dilated cardiomyopathy, 1358
- Cardiac rehabilitation
- new paradigm for resistance exercise guidelines, 281
 - 12-week, heart rate recovery in heart failure patients after, 694
- Cardiac resynchronization therapy
- criteria, prognostic value for worsening heart failure or arrhythmia in idiopathic dilated cardiomyopathy, 393
 - echocardiographic vs clinical parameters in assessment of response to, 260
 - effect on left ventricular reverse remodeling in CHF patients, 876
 - in heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, relation of invasive hemodynamics to Doppler optimization of atrioventricular delay in, 552
 - mechanism of diastolic mitral regurgitation in candidates for, 1611
 - optimal left ventricular pacing site and tissue synchronization imaging in, 1615
 - in patients with CAD or idiopathic dilated cardiomyopathy, mechanism of cardiac output gain from, 1358
 - relation of right ventricular peak systolic pressure to major adverse events in, 1737
 - "response score" to predict hemodynamic and clinical outcomes, 1732
- Cardiac surgery, clinician use of perioperative testing guidelines, effects on outcomes, 118
- Cardiac syndrome X, microvascular function, metabolic syndrome and risk factor status in women with, 1727
- Cardiac tamponade, repetitive yawning in, 1547 (Readers' Comments)
- Cardiac transplantation. *See* Heart transplantation
- Cardiac volume, in persistent atrial fibrillation before and after cardioversion, 1213
- Cardiologists
- adherence to heart failure treatment guidelines, 151 (Readers' Comments)
 - American, health behavior of, 1093
- Cardiology, good books appearing in 2005, 430 (From the Editor)
- Cardiomyopathy
- Fabry's, impact of enzyme replacement therapy on
 - cardiac morphology and function in, 1515
 - late enhancement in, 1515
 - hypertrophic
 - vs hypertensive left ventricular hypertrophy, diastolic septal perforator flow velocities in, 106
 - and left ventricular apical aneurysms, in identical twins, 1109
 - long-term effects of dual-chamber pacing with echocardiographic evaluation of atrioventricular delay, 1769
 - idiopathic dilated
 - circadian variation of endothelial function, 699
 - decreased cerebral blood flow in CHF secondary to, 1365
 - diastolic and systolic duration in children with heart failure due to, 101
 - disease management program vs primary care physician, 253
 - mechanism of cardiac output gain from CRT in patients with, 1358
 - prognostic value of CRT criteria for worsening heart failure and arrhythmia in, 393
 - inflammatory dilated, atorvastatin for heart failure secondary to, 899
 - ischemic or idiopathic dilated
 - IVA and ejection fraction by tissue Doppler for assessment of left ventricular systolic function in, 872
 - relation of invasive hemodynamics to Doppler optimization of atrioventricular delay in CRT in heart failure secondary to, 552
 - ischemic or non-ischemic, effects of head-up tilt on QTc prolongation in, 412
 - nonischemic dilated, with syncope, high risk of ventricular arrhythmias in patients with, 416
 - obstructive hypertrophic
 - predictors of subacute complete heart block after alcohol septal ablation for, 264
 - time course of pressure gradient response after alcohol septal ablation for, 1511

- peripartum, incidence, mortality and risk factors, 1765
- Cardioprotective effect, of vigorous vs moderate intensity aerobic exercise, 141
- Cardiopulmonary exercise testing. *See* Exercise testing
- Cardiopulmonary resuscitation, out-of-hospital, effects of thrombolysis during, 305
- Cardiovascular disease
- and abdominal aortic aneurysm in men, 1123 (Readers' Comments)
 - adverse
 - effects of tadalafil, 1778
 - in osteoporotic postmenopausal women, effects of raloxifene, 520
 - after CABG, prediction by
 - CRP level, 1697
 - soluble intercellular adhesion molecule-1 level, 1697
 - in China
 - effects of fast foods, 1676 (Readers' Comments)
 - effects of increasing serum cholesterol, 1676 (Readers' Comments)
 - effects of rising blood pressure, 1676 (Readers' Comments)
 - as complications of noncardiac surgery after PTCA vs stenting, 1188
 - in fetal heterotaxy syndrome, 720
 - risk factors for, role in cognitive impairment, 1262
 - risk stratification in metabolic syndrome and diabetes, stress SPECT for, 1538
 - in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin on biomarkers of, 646
- Cardioversion
- atrial and ventricular volume and function in persistent atrial fibrillation before and after, 1213
 - of atrial fibrillation, in mitral stenosis after PBMV, one-year outcome, 1045
 - electrical, effect of atorvastatin on atrial fibrillation recurrence after, 1490
- Cardioverter-defibrillator
- eligibility criteria for patients in a specialized heart failure clinic, 882
 - implantable
 - causing arrhythmia, 1255
 - frequency and causes of inappropriate therapies, 1255
 - prevalence and predictors of electrical storm in users of, 389
 - prognostic value of restrictive mitral filling pattern, 676
 - inappropriate shocks of, relation to severity of heart failure symptoms, 544
 - inspection compliance, video-based training program for, 578
 - new paradigm for cardiac rehabilitation resistance exercise guidelines, 281
- Carotid arteries
- intima-media thickness, relation to angiographic severity and extent of CAD, 624
 - stenting, optimal activated clotting time during, 1657
 - stiffness, relation to obesity and metabolic syndrome in children, 528
- Catheter ablation
- for atrial fibrillation
 - left atrial reverse remodeling after, 1220
 - plasma atrial natriuretic peptide and brain natriuretic peptide levels after, 1741
 - implications of proximity of coronary arteries to atrioventricular annulus in young patients, 1752
- Caucasian. *See* European continental ancestry group
- CD40 ligand, soluble
- circulating levels in patients with unstable angina undergoing coronary stenting, impact of clopidogrel, 192
 - in predicting stent restenosis in stable angina, 198
 - in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646
- CD44-v6, expression in smooth muscle cells in postnatal remodeling of ductus arteriosus, 1056
- Cerebral blood flow, decreased in CHF secondary to idiopathic dilated cardiomyopathy, 1365
- Cerebral ischemia
- incidence in atrial fibrillation
 - role of hyperviscosity, 1745
 - role of inflammation, 1745
 - transesophageal echocardiography in detection of cardiac sources of embolism in, 894
 - nonbacterial thrombotic endocarditis in, 894
- CHD. *See* Coronary heart disease
- Chest pain
- acute, frequency and timing of changes in TIMI risk score during ER stay in low-risk patients with, 781
 - troponin-positive, contrast-enhanced MRI in detection of myocardial scar in, 768
 - without preexisting myocardial ischemia, prognostic value of
 - clinical risk score, 633
 - early exercise testing, 633
- CHF. *See* Congestive heart failure
- Children. *See* Pediatrics
- Chinese
- cardiovascular disease in
 - effects of fast foods, 1676 (Readers' Comments)
 - effects of increasing serum cholesterol, 1676 (Readers' Comments)
 - effects of rising blood pressure, 1676 (Readers' Comments)
 - lean, systemic hypertension in
 - impact of body mass index, 839
 - impact of dysglycemia, 839
 - impact of waist-to-hip ratio, 839
- Cholesterol. *See also* High-density lipoproteins; Low-density lipoproteins
- increasing serum levels, effects on cardiovascular disease in China, 1676 (Readers' Comments)
 - influence in CHD development, 372
 - management by targeting lower low-density lipoprotein concentration, 1667 (Editorial)
- Chronic heart failure. *See* Heart failure
- Chronic kidney disease, identification of CAD and left ventricular hypertrophy in asymptomatic patients with
 - B-type natriuretic peptide, 1530
 - N-terminal ProBNP, 1530
- Churg-Strauss syndrome, heart disease in, 1519
- Circadian rhythm, variation
- of endothelial function in idiopathic dilated cardiomyopathy, 699
 - in symptomatic subacute stent thrombosis after bare metal coronary stent implantation, 195
- CKD. *See* Chronic kidney disease
- CK-MB. *See* Creatine kinase-MB
- Clinical trials, nonrandomized, in acute coronary syndrome, drawbacks of, 151 (Readers' Comments)
- Clopidogrel
- and aspirin
 - effects on platelet function, 38
 - long-term use after PCI, 38
 - high vs low loading dose, effects on stable angina patients subjected to PCI, 984
 - impact on circulating levels of soluble CD40 ligand in patients with unstable angina undergoing coronary stenting, 192
- Cobalt-chromium alloy
- stents, safety and efficacy for stable and unstable angina pectoris, 349
- Cocaine
- acute administration in habitual users, causing QT interval prolongation, 1244
 - effects of long-term use on left regional ventricular function, 1085
- Cognitive impairment, role of cardiovascular disease risk factors in, 1262
- Cohn, Lawrence Harvey, 929 (Conversation With the Editor)
- Colesevelam hydrochloride, addition to statins, in hypercholesterolemia
 - effects on high-sensitivity CRP, 1198
 - effects on low-density lipoprotein, 1198
- Collateral circulation, coronary vessel development
 - in diabetes mellitus, effect of statins, 772
 - in patients with total right coronary artery occlusion, relation to metabolic syndrome, 636
 - relation to angiogenic factors, 472
- Complement system proteins, C3 and C4, effects on incidence of atrial fibrillation, 245
- Complete heart block, subacute, after alcohol septal ablation for obstructive hypertrophic cardiomyopathy, predictors of, 264
- Computed tomography. *See also* Optical coherence tomography; Positron emission tomography
- multidetector

- vs intravascular ultrasound, characterization of coronary atherosclerotic plaques, 598
- vs noninvasive coronary angiography, in detection of coronary stenosis before valve surgery, 1506
- for noninvasive evaluation of coronary arteries in asymptomatic patients, 287 (Editorial)
- multidetector row spiral, for detection of significant coronary artery stenosis, 343
- 64-slice cardiovascular, vs coronary angiography, in suspected CAD, 173
- Congenital heart disease
 - after cardiac surgery, incidence and risk factors of heart failure in adolescents and adults with, 1238
 - causing volume and/or pressure overload
 - effects on B-type natriuretic peptides, 1051
 - effects on right ventricular function, 1051
- Congestive heart failure
 - cost-effectiveness of Doppler vs B-type natriuretic peptide in predicting outcomes of, 400
 - effect of CRT on left ventricular reverse remodeling in, 876
 - hospital admissions for, relation to air pollution in seven United States cities, 404
 - impact on non-ST-segment elevation ACS, 1707
 - in Japanese AMI patients, effects of statins, 1165
 - with preserved ejection fraction, gender differences in mortality and morbidity, 1228
 - quality of care in African-Americans admitted at a university teaching hospital, 690
 - relation of endothelial activation, dysfunction, and damage to B-type natriuretic peptide and outcomes, 671
 - secondary to idiopathic dilated cardiomyopathy, decreased cerebral blood flow in, 1365
- Contrast
 - MRI
 - in detection of myocardial scar in troponin-positive chest pain and minimal angiographic CAD, 768
 - vs electrocardiography, in detection of healed MI, 443
 - vs fluorine-18 deoxyglucose PET, in detection of myocardial viability, 452
 - myocardial echocardiography, vs SPECT, to predict AMI transmural, 1718
 - nephropathy induced by, incidence in high-risk patients, effect of rapid intra-arterial infusion of 5% dextrose before coronary angiography, 981
- Contrast echocardiography. *See* Echocardiography
- Coronary angiography
 - extent of CAD, relation to T wave abnormalities, 1124 (Readers' Comments)
 - noninvasive, vs multidetector computed tomography, in detection of coronary stenosis before valve surgery, 1506
 - rapid intra-arterial infusion of 5% dextrose before, 981
 - relation of obesity to CAD severity, 1277
 - vs 64-slice cardiovascular computed tomography, in suspected CAD, 173
- Coronary arteries
 - atherosclerotic plaques, characterization using intravascular ultrasound vs multidetector computed tomography, 598
 - calcified plaque
 - relation to CHD, 1564
 - relation to left ventricular hypertrophy, effect of ethnicity, 1441
 - calcium
 - relation to ankle-brachial index, 181
 - vs stress myocardial perfusion imaging in apparently healthy siblings of individuals with premature CAD, 328
 - infarct-related, effect of medication use prior to STEMI on patency of, 7
 - intravascular ultrasound of, during PCI in ACS, 29
 - late luminal loss as predictor of clinical efficacy of drug-eluting stents in, 603
 - microvascular resistance, during anterior wall AMI and after PCI, 1131
 - minimally diseased, remodeling patterns of, 13
 - multiple unstable plaques, manifested by recurrent STEMI in different territory after fibrinolysis, 947
 - native, vs saphenous vein grafts, intravascular ultrasound of ruptured plaques in, 593
 - noninvasive evaluation with multidetector computed tomography in asymptomatic patients, 287 (Editorial)
 - plaques, assessment by optical coherence tomography, 1172
 - predictors and incidence of restenosis after placement of drug-eluting stents in, 506
 - proximity to atrioventricular annulus in young patients, implications in catheter ablation, 1752
 - unprotected left main, safety and feasibility of various sirolimus-eluting stent implantation techniques for bifurcation stenosis of, 1597
- Coronary arteriography. *See* Coronary angiography
- Coronary artery bypass grafting
 - in the drug-eluting stent era, rescue PCI after, 789
 - effect of statins on risk of atrial fibrillation after, 55, 1266 (Readers' Comments), 1551 (Readers' Comments)
 - new paradigm for cardiac rehabilitation resistance exercise guidelines, 281
 - off-pump, risk and determinants of myocardial injury during, 1482
 - on- vs off-label use of drug-eluting stents in clinical practice, 1478
 - preoperative CRP and soluble intercellular adhesion molecule-1 level for predicting future cardiovascular events after, 1697
 - stress testing after, 810
- Coronary artery disease. *See also* Angina pectoris
 - after orthotopic transplant, usefulness and safety of PCI for, 1192
 - angiographic
 - and long-term mortality, gender differences in relation of renal impairment to, 630
 - relation of ascending aortic pressures to outcomes in patients with, 590 (Readers' Comments)
 - in asymptomatic CKD patients, identification by N-terminal ProBNP and B-type natriuretic peptide, 1530
 - effectiveness of "direct" stenting without balloon predilatation, 316
 - effect of statins on risk of ventricular arrhythmia in, 153 (Readers' Comments)
 - effects of obesity on benefits of atorvastatin or pravastatin, 1553
 - effects of sirolimus- vs paclitaxel-eluting stents on T-cell subset redistribution, 494
 - extent
 - in left bundle branch block with or without type 2 diabetes, 857
 - in relation to T wave abnormalities, 1124 (Readers' Comments)
 - in type 2 diabetes with or without left bundle branch block, 857
 - lipid and lipoprotein biomarkers, 640
 - mechanism of cardiac output gain from CRT in patients with, 1358
 - minimal angiographic, contrast-enhanced MRI in detection of myocardial scar in, 768
 - multivessel, after incomplete revascularization, exercise myocardial scintigraphy, 207
 - normal stress technetium-99m-tetrofosmin SPECT in, 1
 - with or without diabetes mellitus, coronary risk factors and quality of life in patients, 1267
 - ostial de novo left anterior descending coronary artery lesions, treated with drug-eluting stents, early outcome of, 187
 - patients undergoing TAPSE, relation of survival to revascularization vs medical therapy in CAD, 804
 - peripheral arterial disease as marker of risk for abdominal aortic aneurysm in, 1549 (Readers' Comments)
 - prediction using P waves during exercise testing, 1126 (Readers' Comments)
 - premature, coronary calcium vs stress myocardial perfusion imaging in apparently healthy siblings of individuals with, 328
 - prevalence in postmenopausal women with or without osteoporosis or osteopenia, 1427
 - relation of carotid intima-media thickness to angiographic severity and extent of, 624
 - relation of short-term sibutramine therapy to weight loss and improved endothelial function in obese patients with, 1650
 - severity
 - in diabetes mellitus, relation to hemoglobin A_{1c} levels, 968
 - in patients undergoing coronary angiography, relation to obesity, 1277
 - stable, effect of ischemic mitral regurgitation on left ventricular filling pressures in, 1449
 - suspected
 - coronary angiography vs 64-slice cardiovascular computed tomography in, 173
 - estrogen status as prognostic marker in women with, 367
 - wall motion abnormalities, during recovery after acute beta-blocker administration, dobutamine stress echocardiography in observation of, 462

- without coronary vasospasm, vs stable angina and ACS with CAD, serum levels of inflammatory markers in patients with, 1429
- with vs without metabolic syndrome, effect of simvastatin on outcomes of adults with, 1487
- Coronary artery occlusion
- acute, PTCA-induced, relation of ST-segment deviation to myocardial ischemia during, 295
 - chronic total
 - intravascular ultrasound of morphologic features, 1455
 - on- vs off-label use of drug-eluting stents in clinical practice, 1478
 - trends in PCI for, 1691
 - distal vs proximal, prognostic effects on AMI after PCI, 1137
 - total, angiogenic factors in, 472
 - total right, relation of metabolic syndrome to coronary collateral vessel development in patients with, 636
- Coronary artery syndrome, early, in families with frequent MI, relation to metabolic syndrome, 964
- Coronary heart disease
- vs all-cause mortality risk, for electrocardiographic abnormalities in men and women, 309
 - coronary sinus and ascending aorta levels of
 - aldosterone, 1068, 1107 (Editorial)
 - angiotensin II, 1068, 1107 (Editorial)
 - B-type natriuretic peptide, 1068, 1107 (Editorial)
 - development of, influence of lipids, 372
 - impact of over-the-counter statin availability, 851
 - incident, effects of gender and race on prognostic value of electrocardiography, 1176, 1176.e1
 - long-term risk in men
 - apo-B and, 997
 - low-density lipoprotein and, 997
 - placebo effect on
 - laser myocardial revascularization in, 152 (Readers' Comments)
 - therapeutic angiogenesis in, 152 (Readers' Comments)
 - prognostic effects of stress and type D personality, 970
 - relation of depression and fatigue to PAI-1, t-PA antigen and fibrinogen levels in patients with or without, 1287
 - relation of T-wave alternans to regional left ventricular dysfunction and eccentric hypertrophy secondary to, 775
 - relationship of serum high-density lipoprotein levels to mortality in, 466
 - relation to coronary artery calcified plaque, 1564
 - and retinal arteries, 1549 (Readers' Comments)
 - risk assessment in, 765
- Coronary revascularization, normal stress technetium-99m-tetrofosmin SPECT in, 1
- Coronary sinus, of aortic stenosis and CHD patients
- aldosterone levels in, 1068, 1107 (Editorial)
 - angiotensin II levels in, 1068, 1107 (Editorial)
 - B-type natriuretic peptide levels in, 1068, 1107 (Editorial)
- Coronary stenosis
- significant, detection with multidetector row spiral computed tomography, 343
 - before valve surgery, detection with multidetector computed tomography vs non-invasive coronary angiography, 1506
- Coronary thrombus
- of AMI patients
 - fibrin in, 26
 - platelets in, 26
 - TF in, 26
 - vWF in, 26
 - assessment by optical coherence tomography, 1713
- Coronary vasospasm
- induced by ergonovine maleate in vasospastic angina pectoris, intravascular ultrasound of, 1322
 - without CAD, vs stable angina and ACS with CAD, serum levels of inflammatory markers in patients with, 1429
- Cost-effectiveness, of Doppler vs B-type natriuretic peptide in predicting outcomes of CHF, 400
- Counterpulsation
- enhanced external
 - for left ventricular dysfunction, two-year outcomes after, 17
 - for refractory angina pectoris, two-year outcomes after, 17
 - intra-aortic balloon, 1391
- CPR. *See* Cardiopulmonary resuscitation
- C-reactive protein
- after implantation of sirolimus- vs paclitaxel- vs dexamethasone-eluting stents, 1311
 - in atrial fibrillation with or without hypertension, 1346
 - and C3 and C4, effects on incidence of atrial fibrillation, 245
 - high-sensitivity
 - in hypercholesterolemia, efficacy of adding colesvelam hydrochloride to statins, 1198
 - relation to first onset and recurrence of lone atrial fibrillation, 659
 - in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646
 - and nocturnal melatonin levels in STEMI patients, 10
 - for predicting long-term mortality
 - in diabetes mellitus, 167
 - in non-ST-elevation ACS, 167
 - in prediction of left ventricular hypertrophy in systemic hypertension, 249
 - as predictor of aortic valve stenosis progression, 90
 - preoperative level, for predicting future cardiovascular events after CABG, 1697
 - relation to social integration in men and women 70 to 79 years, 1010
 - serum
 - effect of thiazolidinediones on, meta-analysis, 655
 - in patients with coronary vasospasm without CAD vs patients with stable angina and ACS with CAD, 1429
 - relation to heart rate variability in unstable angina, 1702
- Creatine kinase-MB, relation to angiographic adverse events during PCI, 1591
- Creatinine, for predicting long-term mortality
- in diabetes mellitus, 167
 - in non-ST-elevation ACS, 167
- Cross clamping, aortic, during major vascular surgery, effects of statins on renal function after, 1383
- CRP. *See* C-reactive protein
- CRT. *See* Cardiac resynchronization therapy
- Crush technique, long-term outcomes of sirolimus-eluting stent implantation in coronary bifurcation lesions, 1317
- Cypher stents
- overlapping, angiographic patterns of restenosis after implantation of, 499
 - undeployed, failure to retrieve, 502
- Cytokine, improved profile in advanced chronic heart failure, G-CSF for, 681
- Death, effects of morning vs evening atorvastatin intake after first elective PCI, 44
- Depression
- in men with paroxysmal atrial fibrillation, paroxetine for, 1749
 - in patients with or without CHD
 - fibrinogen levels and, 1287
 - PAI-1 levels and, 1287
 - t-PA antigen levels and, 1287
- Desensitization
- rapid, for parents with aspirin sensitivity, 294 (Readers' Comments)
 - reply to a reader's comment, 294
- Dexamethasone-eluting stents, vs sirolimus- and paclitaxel-eluting stents, outcome, restenosis rates and CRP levels, 1311
- Dextrose 5%
- effect on incidence of contrast-induced nephropathy in high-risk patients, 981
 - rapid intra-arterial infusion before coronary angiography, 981
- Diabetes mellitus
- in Asian Indians, relation to lipoprotein lipase polymorphism, 1337
 - coronary risk factors and quality of life in CAD patients with or without, 1267
 - effect of statins on coronary collateral development in, 772
 - heterogeneity of neointimal distribution of in-stent restenosis in, 340
 - insulin sensitization for prevention of MI in, 651
 - prediabetic subjects, relation of iron stores to endothelial dysfunction in, 1550 (Readers' Comments)
 - prediction of long-term mortality by biomarkers, 167
 - racial differences on survival after MI in, 1551 (Readers' Comments)
 - relation of hemoglobin A_{1c} levels to CAD severity in, 968

- stress SPECT for cardiovascular disease risk stratification, 1538
- Diabetes mellitus, type 1
- cardiac function in, 77
 - effect of glycemic control on left ventricular diastolic function in, 71
 - flow mediated dilatation in, 77
 - vs type 2, aspirin resistance in, 567
- Diabetes mellitus, type 2
- effects of rosiglitazone monotherapy vs combination with atorvastatin on biomarkers of cardiovascular disease in, 646
 - with or without left bundle branch block, extent of CAD in, 857
 - relation of metabolic syndrome and insulin resistance to silent myocardial ischemia in, 236
 - vs type 1, aspirin resistance in, 567
- Diastolic function
- in children with heart failure due to idiopathic dilated cardiomyopathy, 101
 - effects of increasing heart rate on Doppler-derived indexes of, 130
 - left ventricular
 - after exercise training in young men, 1089
 - in Behçet's disease, pulse-wave Doppler and Doppler tissue imaging of, 716
 - effects of growth hormone therapy, 1097
 - effects of long-term cocaine use, 1085
 - effects of quinapril, 916
 - in male body builders, effects of androgenic anabolic steroids, 912
 - in non-compacted and compacted segments in isolated left ventricular non-compaction, 738
 - in patients with erectile dysfunction, 1785
 - in predicting future heart failure, Doppler parameters of, 866
 - in pseudoxanthoma elasticum, 1535
 - relation to B-type natriuretic peptide early after AMI, 1146
 - in type 1 diabetes, effect of glycemic control on, 71
 - in mitral regurgitation, mechanism in CRT candidates, 1611
 - septal perforator flow velocities in hypertrophic cardiomyopathy vs hypertensive left ventricular hypertrophy, 106
 - vs systolic function
 - in heart failure patients ≥ 65 years of age, left atrium in, 83
 - as outcome predictor after PCI for AMI, 160
- Diastolic mitral annular velocity, in predicting exercise tolerance in impaired left ventricular systolic function, 1025
- Diet
- tuna or other broiled or baked fish vs fried fish, effects on cardiac structure, function, and hemodynamics, 216
 - vegetarian, effects on cardiovascular autonomic functions in healthy postmenopausal women, 380
- Dietary supplementation, with omega-3 fatty acids, effects on endothelium-dependent vasodilation in chronic heart failure, 547
- Dilated cardiomyopathy. *See* Cardiomyopathy
- Disease management program, vs primary care physician, for idiopathic dilated cardiomyopathy, 253
- Dobutamine stress echocardiography
- heart rate reserve vs 85% of age-predicted maximum heart rate as measure of chronotropic response during, 742
 - in observing wall motion abnormalities during recovery after acute beta-blocker administration, 462
- Doppler
- accuracy in estimation of right atrial pressure in anesthetized, paralyzed, and mechanically ventilated patients, 1654
 - for assessment of left ventricular systolic function in ischemic or idiopathic dilated cardiomyopathy
 - ejection fractions, 872
 - IVA, 872
 - vs B-type natriuretic peptide, cost-effectiveness in predicting outcomes of CHF, 400
 - contribution of decreased atrial function in pathogenesis of neurally mediated syncope, 1017
 - diastolic mitral annular velocity in predicting exercise tolerance in impaired left ventricular systolic function, 1025
 - effects of androgenic anabolic steroids on left ventricular diastolic function in male bodybuilders, 912
 - effects on increasing heart rate on indexes of diastolic function in, 130
 - left ventricular function after exercise training in young men, 1089
 - optimization of atrioventricular delay in CRT, relation to invasive hemodynamics in heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, 552
 - parameters of left ventricular end-diastolic pressure in predicting future heart failure, 866
 - pulse-wave and tissue imaging, of left ventricular function in Behçet's disease, 716
 - right ventricular free wall mechanical delay and dysfunction, 434 (Readers' Comments)
 - transesophageal, for assessment of aortic wall mechanics in Marfan syndrome, 571
 - transthoracic three-dimensional, vs PISA, for assessment of mitral regurgitation, 1630
- Ductus arteriosus
- postnatal remodeling of, CD44-v6 expression in smooth muscle cells, 1056
 - reverse orientation, significance in neonates with POTO for early intervention, 716
- Dysglycemia, impact on systemic hypertension in lean Chinese, 839
- Dyslipidemia, in Asian Indians, relation to lipoprotein lipase polymorphism, 1337
- Dyssynchrony, ventricular, in compensated heart failure, exercise echocardiographic assessment of effects of biventricular and left ventricular pacing, 1622
- Echocardiography. *See also* Doppler; Transesophageal atrial pacing stress echocardiography; Transesophageal echocardiography; Transthoracic echocardiography
- vs clinical parameters, in assessment of response to CRT, 260
 - diagnosis of fetal heterotaxy syndrome, 720
 - dobutamine stress
 - heart rate reserve vs 85% of age-predicted maximum heart rate as measure of chronotropic response during, 742
 - in observing wall motion abnormalities during recovery after acute beta-blocker administration, 462
 - evaluation of optimal atrioventricular delay with dual-chamber pacing, long-term effects on hypertrophic cardiomyopathy, 1769
- exercise
- effects of biventricular and left ventricular pacing on ventricular dyssynchrony in compensated heart failure, 1622
 - safety in patients with abnormal serum potassium concentration, 1247
- heart disease in Churg-Strauss syndrome, 1519
- intracardiac accessory thyroid gland, 926
- left ventricular systolic and diastolic functions in pseudoxanthoma elasticum, 1535
- vs MRI, to predict myocardial function improvement after PCI, 361
- myocardial contrast, vs SPECT, to predict AMI transmural, 1718
- outpatient laboratory, impacts of obesity, 1082
- real-time three-dimensional
- high-resolution, to visualize left ventricular endocardial surface in MI, 1578
 - vs MRI, in assessment of left ventricular mass, 113
- three-dimensional, vs MRI, feasibility and variability in arrhythmogenic right ventricular dysplasia, 703
- transesophageal vs transthoracic, diagnosis of right-to-left atrial shunting, 434 (Readers' Comments)
- EECP. *See* Enhanced external counterpulsation
- Ejection fractions
- depressed, in men, effects of omega-3 fatty acids on
 - heart rate recovery after exercise, 1127
 - heart rate variability, 1127
 - resting heart rate, 1127
- left ventricular
- effect of body mass index, 725
 - estimation using mitral valve EPSS vs MRI, 137
 - in non-compacted and compacted segments in isolated left ventricular non-compaction, 738
 - normal, left ventricular geometry and survival in patients with, 959
 - preserved, outcomes and hospitalization rates in heart failure with and without, 256

- preserved, gender differences in mortality and morbidity in heart failure with, 1228
- by tissue Doppler, for assessment of left ventricular systolic function in ischemic or idiopathic dilated cardiomyopathy, 872
- Elderly
- Caucasians with MI, relation of IGF-I promoter polymorphism to mortality in, 1274
 - community-dwelling individuals, mortality and morbidity in
 - AAC and, 1281
 - AVS and, 1281
 - MAC and, 1281
 - relation of ventricular arrhythmias to all-cause mortality and AMI in apparently healthy subjects, 1351
- Electrical storm. *See* Ventricular fibrillation; Ventricular tachycardia
- Electrocardiography
- abnormalities in men and women, all-cause mortality risk vs CHD for, 309
 - vs contrast-enhanced MRI, in detection of healed MI, 443
 - prevalence and determinants of left ventricular hypertrophy in multiethnic postmenopausal women, 512
 - prognostic value in incident CHD, effects of gender and race, 1176, 1176.e1
 - QT interval prolongation by acute cocaine administration in habitual users, 1244
 - redefinition of diagnostic Q wave criteria for MI, 974
 - relation of T wave abnormalities to extent of CAD, 1124 (Readers' Comments)
 - routine preoperative, prognostic value in noncardiac surgery, 1103
 - ST-segment deviation in, and troponin T, for early risk stratification in non-ST-segment elevation ACS, 334
 - >10 minutes after ER arrival, in non-ST-segment elevation ACS, frequency and consequences, 437
- Electrophysiology, of atrioventricular nodal reentry tachycardia
- effects of slow pathway ablation, 860
 - gender differences in, 384
- Embolism, cardiac sources of, in cancer and cerebral ischemia, detection by transesophageal echocardiography, 894
- Emergency medical services, vs self-transport, mortality in STEMI patients arriving by, 458
- Emergency room
- frequency and consequences of electrocardiogram recording in non-ST-segment elevation ACS patients >10 minutes after arrival, 437
 - stay in low-risk patients with acute chest pain, frequency and timing of changes in TIMI risk score during, 781
- Empathy, and sympathy, clinical usefulness of, 1789 (Editorial)
- EMS. *See* Emergency medical services
- Endocarditis, nonbacterial thrombotic, in cancer and cerebral ischemia, detection by transesophageal echocardiography, 894
- Endocardium, left ventricular, in MI, visualization by high-resolution real-time three-dimensional echocardiography, 1578
- Endothelium
- activation, dysfunction, and damage in CHF, relation to B-type natriuretic peptide and outcomes, 671
 - dysfunction
 - in prediabetic subjects, relation to iron stores, 1550 (Readers' Comments)
 - relation to fasting plasma glucose and hyperinsulinemia, 590 (Readers' Comments)
 - flow-mediated dilation dependent on, metabolic vasodilation in response to, in African-American vs non-African-American patients with chronic heart failure, 685
 - function
 - in hyperlipidemia, effects of atorvastatin, 1002
 - in idiopathic dilated cardiomyopathy, circadian variation of, 699
 - in obese patients with CAD, relation to short-term sibutramine therapy, 1650
 - in patients with erectile dysfunction, 1785
 - in women with cardiac syndrome X, 1727
 - impaired
 - among Hispanic women, 68
 - and increased body mass index, 68
 - vasodilation in chronic heart failure, effects of dietary omega-3 fatty acid supplementation, 547
- Enhanced external counterpulsation
- for left ventricular dysfunction, two-year outcomes after, 17
 - for refractory angina pectoris, two-year outcomes after, 17
- Enzyme replacement therapy, impact on cardiac morphology and function and late enhancement in Fabry's cardiomyopathy, 1515
- E point-to-septal separation, of mitral valve, vs MRI, in estimation of left ventricular ejection fraction, 137
- EPSS. *See* E point-to-septal separation
- Eptifibatid
- effects on inhibition of platelet aggregation in patients undergoing high-risk PCI, 489
 - for thrombocytopenia, 426
- ER. *See* Emergency room
- Erectile dysfunction
- incidence of cardiovascular adverse events in patients taking tadalafil for, 1778
 - left ventricular diastolic function and endothelial function in patients with, 1785
- Ergonovine maleate, in vasospastic angina pectoris, intravascular ultrasound of coronary vasospasm induced by, 1322
- Errata, 154, 592
- Erythrocyte survival, breath carbon monoxide to measure influence of prosthetic heart valves on, 1374
- Esophageal leads, use in determining strategies of pulmonary vein ablation, 1494
- Estrogen status, as prognostic marker in women with suspected CAD, 367
- Ethnicity
- as determinant of electrocardiographic left ventricular hypertrophy in postmenopausal women, 512
 - effect on association of coronary atherosclerosis to left ventricular hypertrophy, 1441
 - effect on prognostic value of electrocardiography in incident CHD, 1176, 1176.e1
 - effects on survival after MI among diabetics, 1551 (Readers' Comments)
 - QT interval and subintervals from women aged 50 to 79 years, 730
- European continental ancestry group
- vs African-Americans, phenotypic predictors of response to simvastatin therapy, 843
 - elderly, relation of IGF-I promoter polymorphism to mortality in MI, 1274
 - men and women, blood pressure stress reactivity and left ventricular mass among, 240
- Exercise
- abnormal heart rate recovery after, relation to left ventricular perfusion and function, 1452
 - extreme endurance running, effects on HRV in healthy trained subjects, 276
 - home-based program, for overweight and obese patients with advanced heart failure, 886
 - in men with healed MI and depressed ejection fraction, effects of omega-3 fatty acids on heart rate recovery after, 1127
 - metabolic vasodilation in response to, in African-American vs non-African-American patients with chronic heart failure, 685
 - vigorous vs moderate intensity, cardioprotective effects of, 141
- Exercise capacity
- after univentricular vs biventricular repair for pulmonary atresia/intact ventricular septum, 1638
 - reduced, relation to biventricular dysfunction in adult patients with operated tetralogy of Fallot, 1377
- Exercise testing
- in CAD prediction, P waves during, 1126 (Readers' Comments)
 - cardiopulmonary
 - in pulmonary hypertension, 123
 - and six-minute walk, 123
 - early, prognostic value in chest pain without preexisting myocardial ischemia, 633
 - echocardiography, effects of biventricular and left ventricular pacing on ventricular dyssynchrony in compensated heart failure, 1622
 - myocardial perfusion, relation of myocardial ischemia to ventricular premature complexes during recovery from, 1570
 - myocardial scintigraphy, in multivessel coronary disease after incomplete revascularization, 207
 - safety in patients with abnormal serum potassium concentration, 1247
 - ventilatory equivalent vs peak oxygen consumption in predicting outcomes of heart failure patients, 1607

- Exercise tolerance
 effects of growth hormone therapy, 1097
 in impaired left ventricular systolic function, prediction by diastolic mitral annular velocity, 1025
- Exercise training
 in cardiac rehabilitation, new paradigm for, 281
 effects on heart rate recovery in heart failure, 694
 effects on left ventricular function in young men, 1089
- Ezetimibe/simvastatin (Vytorin), vs simvastatin monotherapy, muscle safety profiles, 223
- Fabry's disease, impact of enzyme replacement therapy on cardiac morphology and function and late enhancement in cardiomyopathy due to, 1515
- Fast foods, effects on cardiovascular disease in China, 1676 (Readers' Comments)
- Fatigue, relation to PAI-1, t-PA antigen and fibrinogen levels in patients with or without CHD, 1287
- Fenoldopam, intravenous vs intrarenal administration, in patients undergoing cardiac catheterization
 effects on blood pressure, 1076
 effects on renal function, 1076
- ¹⁸F-Fluorodeoxyglucose imaging, PET, vs contrast-enhanced MRI, in detection of myocardial viability, 452
- FFP. *See* Fresh frozen plasma
- Fibrin, in coronary thrombus of AMI patients, 26
- Fibrinogen, relation to depression and fatigue in patients with or without CHD, 1287
- Fibrinolysis
 for acute pulmonary embolism, predictors of hemorrhage after, 127
 vs primary coronary angioplasty, ST-segment elevation for predicting outcomes after, 611
 recurrent STEMI in different territory after, as manifestation of multiple unstable coronary arterial plaques, 947
- Fibroblast growth factor
 relation to development of collateral vessels, angiographic evaluation, 472
 in stable angina pectoris and total coronary artery occlusion, 472
- Filling pressures, left ventricular, in stable CAD, effect of ischemic mitral regurgitation, 1449
- Flow-mediated dilatation
 of brachial artery
 circadian variation in idiopathic dilated cardiomyopathy, 699
 in hyperlipidemia, effects of atorvastatin, 1002
 and cardiac function, 77
 endothelium-dependent, metabolic vasodilation in response to, in African-American vs non-African-American patients with chronic heart failure, 685
 forearm, in healthy volunteers, relation to plasma insulin levels, 1250
 in type 1 diabetes, 77
- Flow velocity reserve, coronary, during anterior wall AMI and after PCI, 1131
- Fluid restriction, vs tolvaptan, for hyponatremia, 1064
- Fontan repair, vs outflow tract reconstruction, for pulmonary atresia/intact ventricular septum, exercise capacity after, 1638
- Forearm, flow-mediated dilatation in healthy volunteers, relation to plasma insulin levels, 1250
- Fractional shortening, relation to pericardial effusion, 910
- Fresh frozen plasma, efficacy and safety for warfarin-related over-anticoagulation in patients with mechanical heart valves, 409
- Functional status, in chronic severe mitral regurgitation, plasma N-terminal ProBNP as marker of, 1498
- G-CSF. *See* Granulocyte-colony stimulating factor
- Gender differences. *See also* Men; Women
 effect on prognostic value of electrocardiography in incident CHD, 1176, 1176.e1
 in electrophysiologic parameters in atrioventricular nodal reentry tachycardia, 384
 in morbidity and mortality in heart failure with preserved ejection fraction, 1228
 in relation of renal impairment to angiographic CAD and long-term mortality, 630
 of revascularization in AMI, 1722
- Glomerular filtration rate, estimated, effect of short-term rosuvastatin treatment, 1602
- Glucose
 blood, relation to long-term mortality in ACS, 1573
 fasting plasma levels, relation to endothelial dysfunction, 590 (Readers' Comments)
 plasma, for predicting in-hospital outcomes after AMI, 1558
- Glycemic control, effect on left ventricular diastolic function in type 1 diabetes, 71
- Glycoprotein IIb/IIIa receptor antagonists
 effects on inhibition of platelet aggregation in patients undergoing high-risk PCI, 489
 planned vs provisional use in smokers undergoing PCI, 1679
 for thrombocytopenia, 426
- GPRAs. *See* Glycoprotein IIb/IIIa receptor antagonists
- Granulocyte-colony stimulating factor, safety and effectiveness in advanced chronic heart failure
 for improving cytokine profile, 681
 for mobilizing hematopoietic stem cells, 681
- Growth hormone
 effects on aortic distensibility and dimensions in Turner syndrome, 1644
 effects on cardiac morphology and function, 1097
- Growth hormone deficiency, effects of growth hormone therapy on cardiac morphology and function in, 1097
- Harrison, Donald Carey, 1399 (Conversation With the Editor)
- Head-up tilt, effects on QTc prolongation in ischemic or non-ischemic cardiomyopathy, 412
- Health behavior, of American cardiologists, 1093
- Health care delivery, patient and physician perceptions on quality of, 1073
- Heart-assist devices, Impella Recover LP 2.5, for left ventricular support in elective high-risk PCI, feasibility and safety, 990
- Heart conduction system, disturbances in, after transcatheter closure of muscular ventricular septal defects using Amplatzer occluder, 558
- Heart disease. *See also* Congenital heart disease
 advanced, transendocardial autologous bone marrow transplantation for, 823
 in Churg-Strauss syndrome, 1519
- Heart failure
 advanced
 home-based exercise program for overweight and obese patients with, 886
 relation of loop diuretic dose to mortality in, 1759
 safety and tolerability of serial home infusions of nesiritide for, 1370
 advanced chronic, safety and effectiveness of G-CSF for
 improving cytokine profile, 681
 mobilizing hematopoietic stem cell, 681
 after 12-week cardiac rehabilitation, heart rate recovery in patients, 694
 cardiologists' adherence to treatment guidelines for, 151 (Readers' Comments)
 chronic
 effects of dietary omega-3 fatty acid supplementation on endothelium-dependent vasodilation in, 547
 metabolic vasodilation in response to exercise, ischemia, and endothelium-dependent flow-mediated dilation in African-Americans vs non-African-Americans with, 685
 compensated, exercise echocardiographic assessment of effects of biventricular and left ventricular pacing to ventricular dyssynchrony in, 1622
 diastolic vs systolic, in persons ≥ 65 years of age, left atrium in, 83
 due to idiopathic dilated cardiomyopathy, in children with
 diastolic duration in, 101
 systolic duration in, 101
 future events of, prediction by Doppler parameters of left ventricular end-diastolic pressure, 866
 implantable cardioverter defibrillator eligibility criteria for patients in a specialized clinic, 882
 incidence and risk factors in adolescents and adults with congenital heart disease after cardiac surgery, 1238
 prediction by aortic root dimension in persons ≥ 65 years of age, 270
 secondary to inflammatory dilated cardiomyopathy and elevated low-density lipoprotein cholesterol levels, atorvastatin for, 899

- secondary to ischemic or idiopathic dilated cardiomyopathy, relation of invasive hemodynamics to Doppler optimization of atrioventricular delay in CRT in, 552
- severity of symptoms, relation to inappropriate defibrillator shocks, 544
- systolic, prognostic value of restrictive mitral filling pattern, 676
- ventilatory equivalent vs peak oxygen exercise consumption in predicting outcomes of patients with, 1607
- with and without intraventricular conduction defect, outcomes and hospitalization rates, 256
- with and without preserved left ventricular ejection fraction, outcomes and hospitalization rates, 256
- worsening, in idiopathic dilated cardiomyopathy, prognostic value of CRT criteria for, 393
- Heart rate**
- increasing, effects on Doppler-derived indexes of diastolic function, 130
- resting, in men with healed MI and depressed ejection fraction, effects of omega-3 fatty acids, 1127
- Heart rate recovery**
- after exercise, in men with healed MI and depressed ejection fraction, effects of omega-3 fatty acids, 1127
- in heart failure patients after 12-week cardiac rehabilitation, 694
- postexercise abnormal, relation to left ventricular perfusion and function, 1452
- Heart rate reserve**, vs 85% of age-predicted maximum heart rate, as measure of chronotropic response during dobutamine stress echocardiography, 742
- Heart rate variability**
- in healthy
- postmenopausal women, effects of vegetarian diet, 380
- trained subjects, effects of extreme endurance running, 276
- in men with healed MI and depressed ejection fraction, effects of omega-3 fatty acids, 1127
- relation to serum CRP in unstable angina, 1702
- Heart septal defects**
- atrial, diagnosis with transesophageal vs transthoracic echocardiography, 434 (Readers' Comments)
- ventricular, transcatheter closure using Amplatzer occluder, cardiac conduction disturbances after, 558
- ventricular arrhythmias after, 558
- Heart transplantation**
- effects of vitamin C to coronary hypersensitivity to serotonin in recipients, 561
- orthotopic, usefulness and safety of PCI for vasculopathy after, 1192
- Hematopoietic stem cell**, mobilization in advanced chronic heart failure, G-CSF for, 681
- Hemodynamics**
- cardiac, effects of tuna or other broiled or baked fish vs fried fish, 216
- invasive, relation to Doppler optimization of atrioventricular delay in CRT in heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, 552
- outcome after CRT, "response score" for prediction of, 1732
- Hemoglobin A_{1c}**, relation to CAD severity in diabetes mellitus, 968
- Hemorrhage**, after fibrinolysis for acute pulmonary embolism, predictors of, 127
- Hemostasis**, manual pressure, after transfemoral coronary angiography, effectiveness in patients on warfarin anticoagulation, 485
- Hepatocyte growth factor**
- relation to development of collateral vessels, angiographic evaluation, 472
- in stable angina pectoris and total coronary artery occlusion, 472
- Heterotaxy syndrome**, fetal, cardiovascular disease, diagnostic accuracy and outcome in, 720
- High-density lipoproteins**
- influence in CHD development, 372
- phenotypic predictors of response to simvastatin therapy among African-Americans vs Caucasians, 843
- serum, relation to cardiac mortality in CHD, 466
- Hispanic Americans**, women
- impaired endothelial function in, 68
- increased body mass index in, 68
- Histology**, myocardial wall of left atrium, relation to pulmonary vein incorporation, 662
- History**, sphygmograph in America, 580
- Hospital costs, in treating complications of PCI, 322
- Hospitalization, length of, in major vascular surgery patients, influence of cardiac risk factors and medication, 1423
- HRV. *See* Heart rate variability
- Hypercholesterolemia**
- African-Americans, efficacy and safety of rosuvastatin vs atorvastatin for, 229
- efficacy of adding colesvelam hydrochloride to statins, 1198
- Hyperemia, reactive, after PCI, effect on myocardial tissue reperfusion in AMI, 617
- Hyperinsulinemia, relation to endothelial dysfunction, 590 (Readers' Comments)
- Hyperlipidemia, effects of atorvastatin on endothelial function in, 1002
- Hypertension**. *See also* Pulmonary hypertension
- CRP in atrial fibrillation with or without, 1346
- with left ventricular hypertrophy, vs hypertrophic cardiomyopathy, diastolic septal perforator flow velocities in, 106
- systemic
- CRP and B-type natriuretic peptide in prediction of left ventricular hypertrophy in, 249
- in lean Chinese, impact of dysglycemia, body mass index and waist-to-hip ratio on, 839
- Hypertrophic cardiomyopathy**. *See* Cardiomyopathy
- Hyperviscosity**, role in incidence of cerebral ischemia in atrial fibrillation, 1745
- Hyponatremia**, tolvaptan vs fluid restriction, 1064
- Idiopathic dilated cardiomyopathy**. *See* Cardiomyopathy
- IGF-I**. *See* Insulin-like growth factor-I
- IL-6**. *See* Interleukin-6
- Impella Recover LP 2.5 system**, for left ventricular support in elective high-risk PCI, feasibility and safety, 990
- Infant**, newborn. *See also* Pediatrics
- significance of reverse orientation of ductus arteriosus for early intervention of POTO, 716
- Inflammation**
- relation to risk factors of valvular calcification, 1502
- role in incidence of cerebral ischemia in atrial fibrillation, 1745
- Inflammatory dilated cardiomyopathy**. *See* Cardiomyopathy
- Insulin**, plasma levels, relation to forearm flow-mediated dilatation in healthy volunteers, 1250
- Insulin-like growth factor-I**
- promoter polymorphism of, relation to mortality in elderly Caucasians with MI, 1274
- serum levels in subjects >65 years of age, relation to ACE inhibitor treatment, 1525
- Insulin resistance**, relation to silent myocardial ischemia in type 2 diabetes mellitus, 236
- Insulin sensitization**, for prevention of MI in diabetes mellitus, 651
- Intact ventricular septum**, exercise capacity after univentricular vs biventricular repair for, 1638
- Intercellular adhesion molecule-1**, soluble
- preoperative level, for predicting future cardiovascular events after CABG, 1697
- serum levels in patients with coronary vasospasm without CAD vs patients with stable angina and ACS with CAD, 1429
- Interleukin-6**
- relation to social integration in men and women 70 to 79 years, 1010
- serum levels in patients with coronary vasospasm without CAD vs patients with stable angina and ACS with CAD, 1429
- Interleukins**
- for predicting long-term mortality
- in diabetes mellitus, 167
- in non-ST-elevation ACS, 167
- Intra-aortic balloon counterpulsation**, 1391
- Intracardiac accessory thyroid gland**, 926
- Intracoronary brachytherapy**, for in-stent restenosis, serial angiography after, 21
- Intravascular ultrasound**
- characterization of the "black hole" phenomenon after sirolimus-eluting stent implantation, 203
- comparison of ruptured plaques in native coronary arteries vs saphenous vein grafts, 593

- of coronary vasospasm induced by ergonovine maleate in vasospastic angina pectoris, 1322
- heterogeneity of neointimal distribution of in-stent restenosis in diabetes mellitus, 340
- morphologic features of chronic total coronary occlusions, 1455
- morphology of aorto-ostial vs shaft saphenous vein graft lesions, 1463
- vs multidetector computed tomography, characterization of coronary atherosclerotic plaques, 598
- no-reflow phenomenon during stent implantation in AMI, 301
- during PCI in ACS, coronary arterial finding by, 29
- of restenosis after drug-eluting stent implantation vs brachytherapy, 1292
- Intraventricular conduction defect, outcomes and hospitalization rates in heart failure with and without, 256
- Iron stores, relation to endothelial dysfunction in prediabetic subjects, 1550 (Readers' Comments)
- Ischemia. *See also* Cerebral ischemia; Myocardial ischemia
 - metabolic vasodilation in response to, in African-American vs non-African-American patients with chronic heart failure, 685
 - relation to angiographic adverse events during PCI, 1591
- Ischemic cardiomyopathy. *See* Cardiomyopathy
- Ischemic stroke. *See* Stroke
- Isovolumic acceleration rate, by tissue Doppler, for assessment of left ventricular systolic function in ischemic or idiopathic dilated cardiomyopathy, 872
- IVA. *See* Isovolumic acceleration rate

- Japanese, effects of statins on CHF and myocardial ischemia in AMI patients, 1165

- Kidney, effects of atorvastatin 80 mg vs 10 mg on, 65
- Kidney disease, chronic, CAD and left ventricular hypertrophy in asymptomatic patients with
 - B-type natriuretic peptide and, 1530
 - N-terminal ProBNP and, 1530
- Kidney function
 - after aortic cross clamping during major vascular surgery, effects of statins, 1383
 - effects of intravenous vs intrarenal fenoldopam in patients undergoing cardiac catheterization, 1076
 - effects of statins, 748

- Lead aVR. *See* Electrocardiography
- Left anterior descending artery, ostial de novo lesions, treated with drug-eluting stents, early outcome of, 187
- Left atrium
 - appendage thrombus, detection by transesophageal echocardiography 421
 - histology of myocardial wall of, relation to pulmonary vein incorporation, 662
 - remodeling
 - effects of quinapril, 916
 - reverse, after catheter ablation for atrial fibrillation, 1220
 - structure and function, effects of tuna or other broiled or baked fish vs fried fish, 216
 - in systolic vs diastolic heart failure in persons ≥ 65 years of age, 83
- Left bundle branch block. *See* Bundle branch block
- Left ventricle
 - apical aneurysms, and hypertrophic cardiomyopathy in identical twins, 1109
 - cavity size and ejection fraction, effect of body mass index, 725
 - dysfunction
 - in adult patients with operated tetralogy of Fallot, relation to elevated NT-pro-BNP and reduced exercise capacity, 1377
 - secondary to CHD, relation to T-wave alternans, 775
 - endocardial surface in MI, visualization by high-resolution real-time three-dimensional echocardiography, 1578
 - end-systolic remodeling, in organic and functional mitral regurgitation, relation to B-type natriuretic peptide activation, 1029
 - filling pressures, in stable CAD, effect of ischemic mitral regurgitation, 1449
 - geometry, in patients with normal left ventricular ejection fraction, 959
 - involvement in Churg-Strauss syndrome, 1519
 - morphology, in Fabry's cardiomyopathy, impact of enzyme replacement therapy, 1515
 - optimal pacing site, and tissue synchronization imaging in CRT, 1615
 - pacing, and biventricular pacing, effects on ventricular dyssynchrony in compensated heart failure, exercise echocardiographic assessment, 1622
 - remodeling
 - during long-term right ventricular pacing after His bundle ablation, 1223
 - reverse, in CHF patients, effect of CRT, 876
 - support with Impella Recover LP 2.5, feasibility and safety in elective high-risk PCI, 990
- Left ventricular function
 - after exercise training in young men, 1089
 - in Behçet's disease, pulse-wave Doppler and Doppler tissue imaging of, 716
 - diastolic
 - effects of quinapril, 916
 - in male body builders, effects of androgenic anabolic steroids, 912
 - in patients with erectile dysfunction, 1785
 - relation to B-type natriuretic peptide early after AMI, 1146
 - diastolic and systolic, in pseudoxanthoma elasticum, 1535
 - diastolic vs systolic, as outcome predictor after PCI for AMI, 160
 - effects of growth hormone therapy, 1097
 - effects of long-term cocaine use, 1085
 - ejection fraction
 - estimation using mitral valve EPSS vs MRI, 137
 - normal, left ventricular geometry and survival in patients with, 959
 - preserved, outcomes and hospitalization rates in heart failure with and without, 256
 - end-diastolic pressure, in predicting future heart failure, Doppler parameters of, 866
 - in Fabry's cardiomyopathy, impact of enzyme replacement therapy, 1515
 - in non-compacted and compacted segments in isolated left ventricular noncompaction, 738
 - relation to postexercise abnormal heart rate recovery, 1452
 - and structure, effects of tuna or other broiled or baked fish vs fried fish, 216
 - systolic
 - impaired, prediction of exercise tolerance by diastolic mitral annular velocity, 1025
 - in ischemic or idiopathic dilated cardiomyopathy, tissue Doppler assessment of ejection fractions and IVA, 872
 - relation to pericardial effusion, 910
- Left ventricular hypertrophy
 - in asymptomatic CKD patients, identification by N-terminal ProBNP and B-type natriuretic peptide, 1530
 - electrocardiographic, in multiethnic postmenopausal women, prevalence and determinants of, 512
 - hypertensive, vs hypertrophic cardiomyopathy, diastolic septal perforator flow velocities in, 106
 - relation to coronary atherosclerosis, effect of ethnicity, 1441
 - secondary to CHD, relation to T-wave alternans, 775
 - in systemic hypertension, prediction with
 - B-type natriuretic peptides, 249
 - CRP, 249
- Left ventricular mass
 - and blood pressure stress reactivity among Caucasian and Africa-American men and women, 240
 - effects of growth hormone therapy, 1097
 - real-time three-dimensional echocardiography vs MRI in assessment of, 113
- Left ventricular noncompaction, isolated, left ventricular function in non-compacted and compacted segments in, 738
- Leukocyte count. *See* White blood cell count
- Life style, of American cardiologists, 1093
- Lipids
 - biomarkers for CAD, 640
 - in hyperlipidemia, effects of atorvastatin, 1002
 - influence on CHD development, 372
 - phenotypic predictors of response to simvastatin therapy among African-Americans vs Caucasians, 843
 - in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646

- Lipoprotein lipase, polymorphism of, relation to dyslipidemia and diabetes in Asian Indians, 1337
- Lipoproteins. *See also* High-density lipoproteins; Low-density lipoproteins biomarkers for CAD, 640
phenotypic predictors of response to simvastatin therapy among African-Americans vs Caucasians, 843
- Liver, effects of atorvastatin 80 mg vs 10 mg on, 65
- Loop diuretics, dose, relation to mortality in advanced heart failure, 1759
- Loscalzo, Joseph, 1110 (Conversation With the Editor)
- Lovastatin, impact of over-the-counter availability on CHD, 851
- Low-density lipoproteins
effect on atherogenic particle number in predicting risk of fatal MI, 943
elevated levels, atorvastatin for heart failure secondary to, 899
in hypercholesterolemia, efficacy of adding colesvelam hydrochloride to statins, 1198
influence on CHD development, 372
and long-term risk of CHD in men, 997
lower concentration as target of cholesterol management, 1667 (Editorial)
particle size, phenotype B and particle concentrations in Asian Indians, prediction by triglyceride/high-density lipoprotein ratio, 1007
in patients on statin therapy, effects of plant stanol on lowering of, 376
phenotypic predictors of response to simvastatin therapy among African-Americans vs Caucasians, 843
- MAC. *See* Mitral annular calcification
- Magnetic resonance imaging
atrial and ventricular volume and function in persistent atrial fibrillation before and after cardioversion, 1213
contrast
in detection of myocardial scar in troponin-positive chest pain and minimal angiographic CAD, 768
vs electrocardiography, in detection of healed MI, 443
vs fluorine-18 deoxyglucose PET, in detection of myocardial viability, 452
in diagnosis of arrhythmogenic right ventricular dysplasia in children, 1232
vs echocardiography, to predict myocardial function improvement after PCI, 361
effects of long-term cocaine use on left ventricular function, 1085
harmonic phase, for validation of two-dimensional strain and tissue based strain, 1661
intracardiac accessory thyroid gland, 926
vs mitral valve EPSS, in estimation of left ventricular ejection fraction, 137
vs real-time three-dimensional echocardiography, in assessment of left ventricular mass, 113
relation of B-type natriuretic peptide to left ventricular diastolic function and extent of myocardial scar early after AMI, 1146
right ventricular function, effects of volume and/or pressure overload secondary to congenital heart disease, 1051
vs three-dimensional echocardiography, feasibility and variability in arrhythmogenic right ventricular dysplasia, 703
- Males. *See* Men
- Marfan syndrome
aortic root distensibility vs diameter as risk factor in, 1422 (Readers' Comments)
assessment of aortic wall mechanics by transesophageal tissue Doppler echocardiography, 571
- Matrix metalloproteinases
MMP-9
in AMI vs unstable angina pectoris vs stable angina pectoris, 175
in type 2 diabetes, effects of rosiglitazone monotherapy vs combination with atorvastatin, 646
and TIMPs, selective induction in atrial and ventricular myocardium in patients with atrial fibrillation, 532
- Mechanical ventilation, accuracy of Doppler estimation of right atrial pressure in anesthetized and paralyzed patients during, 1654
- Melatonin, nocturnal levels
and CRP in STEMI patients, 10
prognostic value in STEMI, 1162
- Membrane attack complex, expression in failing human myocardium, 1626
- Men
all-cause mortality risk vs CHD for electrocardiographic abnormalities in, 309
cardiovascular disease and abdominal aortic aneurysm in, 1123 (Readers' Comments)
Caucasian and Africa-American, blood pressure stress reactivity and left ventricular mass among, 240
depressed, with paroxysmal atrial fibrillation, paroxetine for, 1749
effects of androgenic anabolic steroids on left ventricular diastolic function in bodybuilders, 912
with healed MI and depressed ejection fractions, effects of omega-3 fatty acids on
heart rate recovery after exercise, 1127
heart rate variability, 1127
resting heart rate, 1127
long-term risk of CHD in
apo-B and, 997
low-density lipoprotein and, 997
and women, 70 to 79 years, relation of social integration to CRP and IL-6 levels, 1010
young, left ventricular function after exercise training in, 1089
- Meta-analysis
effect of thiazolidinediones on serum CRP levels, 655
of rescue PCI after failed fibrinolysis, 1685
- Metabolic syndrome
benefits of niacin in healed MI patients with vs without, 477
effect of simvastatin on cardiovascular outcomes of adults with vs without, 1487
effects of statins on platelet expression of PAR-1 thrombin receptor in patients with, 1332
and insulin resistance, relation to silent myocardial ischemia in type 2 diabetes mellitus, 236
and obesity, relation to carotid artery stiffness in children, 528
relation to body mass index and leukocyte count, 835
relation to coronary collateral vessel development in patients with total right coronary artery occlusion, 636
relation to early CAD in families with frequent MI, 964
stress SPECT for cardiovascular disease risk stratification, 1538
in women with cardiac syndrome X, 1727
- MI. *See* Myocardial infarction
- Middle-aged, relation of ventricular arrhythmias to all-cause mortality and AMI in apparently healthy subjects, 1351
- Military, young female recruits, sudden death in, 1756
- Mitral annular calcification, in community-dwelling elderly individuals
relation to morbidity, 1281
relation to mortality, 1281
- Mitral annular velocity, diastolic, in predicting exercise tolerance in impaired left ventricular systolic function, 1025
- Mitral regurgitation
assessment using PISA vs transthoracic three-dimensional Doppler echocardiography, 1630
chronic severe, plasma N-terminal ProBNP as marker of functional capacity and left ventricular remodeling in, 1498
diastolic, mechanism in CRT candidates, 1611
ischemic, effect on left ventricular filling pressures in stable CAD, 1449
organic and functional, relation of B-type natriuretic peptide activation to left ventricular end-systolic remodeling in, 1029
secondary to mitral valve prolapse, frequency of atrial fibrillation in, 1039
- Mitral valve
calcification, relation of inflammation to risk factors of, 1502
EPSS, vs MRI, in estimation of left ventricular ejection fraction, 137
replacement for rheumatic mitral stenosis, atrial fibrillation prior to, 1035
restrictive filling pattern, prognostic value in systolic heart failure and implantable cardioverter-defibrillator, 676
Starr-Edwards model 6120 mechanical prosthesis, 756
tumor excision vs valve replacement for papillary fibroelastoma of, 759
- Mitral valve prolapse, frequency of atrial fibrillation in mitral regurgitation secondary to, 1039
- Mitral valve stenosis
after PBMV, one-year outcome of cardioversion of atrial fibrillation in, 1045
rheumatic, atrial fibrillation before isolated mitral valve replacement for, 1035
- MMP. *See* Matrix metalloproteinases

- Monocyte chemoattractant protein-1, serum levels in patients with coronary vasospasm without CAD vs patients with stable angina and ACS with CAD, 1429
- Mononuclear cells, autologous bone marrow–derived, intramyocardial injection for severe angina pectoris, 1326
for stress-induced myocardial ischemia, 1326
- Morbidity
in community-dwelling elderly individuals, relation to MAC, AVS and AAC, 1281
in heart failure with preserved ejection fraction, gender differences, 1228
- Mortality. *See also* Sudden death
in abdominal aortic aneurysm patients treated with and without statins, 279
in advanced heart failure, relation to loop diuretic dose, 1759
all-cause
in apparently healthy subjects aged ≥ 55 years, relation to ventricular arrhythmias, 1351
vs CHD, for electrocardiographic abnormalities in men and women, 309
in AMI patients ≥ 65 years of age, relation to statin initiation after hospital discharge, 155
cardiovascular and all-cause, prediction by aortic root dimension in persons ≥ 65 years of age, 270
in community-dwelling elderly individuals
relation to AAC, 1281
relation to AVS, 1281
relation to MAC, 1281
due to pulmonary embolism in ischemic stroke, 1776
in elderly Caucasians with MI, relation to IGF-I promoter polymorphism, 1274
in heart failure with preserved ejection fraction, gender differences, 1228
in-hospital, after balloon angioplasty vs stent implantation for AMI, 977
long-term
after PCI, prediction by neutrophil/lymphocyte ratio, 993
and angiographic CAD, gender differences in relation of renal impairment to, 630
in diabetes mellitus, prediction by biomarkers, 167
in non–ST-elevation ACS, prediction by biomarkers, 167
relation to blood glucose in ACS, 1573
long-term outcomes of PTCA with vs without stenting, 1467
in multivessel vs single-vessel PCI, 1585
one-year, prediction among 30-day STEMI survivors after primary PCI, 817
in peripartum cardiomyopathy, 1765
prediction after AMI
QRS wave, 353
T wave, 353
relation to revascularization vs medical therapy in CAD patients undergoing TAPSE, 804
relation to serum high-density lipoprotein levels in CHD, 466
in STEMI patients arriving by EMS vs self-transport, 458
thirty-day, in AMI, prediction by routine admission laboratory tests, 1435
in users of implantable cardioverter-defibrillators, 1255
- MRI. *See* Magnetic resonance imaging
- Multidetector computed tomography. *See* Computed tomography
- Multislice computed tomography. *See* Computed tomography
- Muscle, safety profiles of Vytorin vs simvastatin monotherapy, 223
- Musculoskeletal system, effects of atorvastatin 80 mg vs 10 mg on, 64
- Myocardial function, improvement after PCI, prediction using MRI vs echocardiography, 361
- Myocardial infarction
acute. *See* Acute myocardial infarction
after multivessel vs single-vessel PCI, 1585
among diabetics, racial differences on survival after, 1551 (Readers' Comments)
angina pectoris before, effect on reperfusion arrhythmias, 1157
effects of morning vs evening atorvastatin intake after first elective PCI, 44
fatal, risk prediction by atherogenic particle size
effects of apo-A-I, 943
effects of apo-B, 943
effects of low-density lipoprotein, 943
frequent, relation of metabolic syndrome to early CAD in families with, 964
healed
detection by contrast-enhanced MRI vs electrocardiography, 443
in men, effects of omega-3 fatty acids on heart rate recovery after exercise, HRV, and resting heart rate, 1127
with vs without metabolic syndrome, benefits of niacin in, 477
high-resolution real-time three-dimensional echocardiography to visualize left ventricular endocardial surface in, 1578
long-term outcomes of PTCA with vs without stenting, 1467
new paradigm for cardiac rehabilitation resistance exercise guidelines, 281
normal stress technetium-99m–tetrofosmin SPECT in, 1
prevention in diabetes mellitus, insulin sensitization for, 651
redefinition of diagnostic Q wave criteria for, 974
relation of IGF-I promoter polymorphism to mortality in elderly Caucasians with, 1274
rescue PCI after CABG in the drug-eluting stent era, 789
ST-segment elevation. *See* ST-segment elevation myocardial infarction
- Myocardial injury, during off-pump CABG, risk and determinants of, 1482
- Myocardial ischemia
in Japanese AMI patients, effects of statins, 1165
preexisting, prognostic value of early exercise testing and clinical risk score in chest pain without, 633
during PTCA-induced acute coronary occlusion, relation to ST-segment deviation, 295
relation to ventricular premature complexes during recovery from myocardial perfusion exercise stress test, 1570
rescue PCI after CABG in the drug-eluting stent era, 789
silent, in type 2 diabetes mellitus, relation to metabolic syndrome and insulin resistance, 236
stress-induced, intramyocardial injection of autologous bone marrow–derived mononuclear cells for, 1326
transendocardial autologous bone marrow transplantation for, 823
- Myocardial perfusion
after MI and/or coronary revascularization, 1
exercise stress test, relation of myocardial ischemia to ventricular premature complexes during recovery from, 1570
relation to postexercise abnormal heart rate recovery, 1452
stress
vs coronary calcium, in apparently healthy siblings of individuals with premature CAD, 328
technetium-99m–tetrofosmin SPECT in, 1
- Myocardial scar
detection by contrast-enhanced MRI
in minimal angiographic CAD, 768
in troponin-positive chest pain, 768
extent of, relation to B-type natriuretic peptide early after AMI, 1146
- Myocardial viability, determination by contrast-enhanced MRI vs fluorine-18 deoxyglucose PET, 452
- Myocardium
atrial and ventricular, in patients with atrial fibrillation, selective induction of TIMPs and MMP in, 532
failing human, expression of membrane attack complex in, 1626
of left atrium, histology of, relation to pulmonary vein incorporation, 662
- Naki, Hamilton, 435 (Readers' Comments)
- Natriuretic peptides
atrial and brain, plasma levels after radiofrequency catheter ablation of atrial fibrillation, 1741
B-type. *See also* N-Terminal ProBNP
activation of, relation to left ventricular end-systolic remodeling in organic and functional mitral regurgitation, 1029
bedside testing for, to identify right ventricular dysfunction and outcome in acute pulmonary embolism, 1386
vs Doppler, cost-effectiveness in predicting outcomes of CHF, 400
effects of volume and/or pressure overload secondary to congenital heart disease on, 1051
to identify CAD and left ventricular hypertrophy in asymptomatic CKD patients, 1530
levels in coronary sinus and ascending aorta of aortic stenosis and CHD patients, 1068, 1107 (Editorial)
in prediction of left ventricular hypertrophy in systemic hypertension, 249

- relation to left ventricular diastolic function and extent of myocardial scar early after AMI, 1146
- relation to outcomes and endothelial activation, dysfunction, and damage in CHF, 671
- Neointima, heterogeneity of, in in-stent restenosis in diabetes mellitus patients, 340
- Nephropathy, contrast-induced, incidence in high-risk patients, effect of rapid intra-arterial infusion of 5% dextrose before coronary angiography on, 981
- Nesiritide, serial home infusions, safety and tolerability for advanced heart failure, 1370
- Neutrophil/lymphocyte ratio, in prediction of long-term mortality after PCI, 993
- Niacin, benefits in healed MI patients with vs without metabolic syndrome, 477
- N-Terminal ProBNP
- on admission of AMI patients, relation to scintigraphic infarct size, reperfusion efficacy and prognosis, 1151
 - elevated, relation to biventricular dysfunction in adult patients with operated tetralogy of Fallot, 1377
 - to identify CAD and left ventricular hypertrophy in asymptomatic CKD patients, 1530
 - plasma, as marker of functional capacity and left ventricular remodeling in chronic severe mitral regurgitation, 1498
 - for predicting long-term mortality
 - in diabetes mellitus, 167
 - in non-ST-elevation ACS, 167
 - for predicting long-term outcome after PCI in patients with normal troponin T levels, 830
- Obesity
- with advanced heart failure, home-based exercise program for, 886
 - with CAD, relation of short-term sibutramine therapy to improved endothelial function, 1650
 - weight loss, 1650
 - effects on benefits of atorvastatin or pravastatin in CAD, 1553
 - impact on outpatient echocardiography laboratory, 1082
 - and leukocyte count, relation to metabolic syndrome, 835
 - and metabolic syndrome, relation to carotid artery stiffness in children, 528
 - relation to CAD severity in patients undergoing coronary angiography, 1277
 - role in cognitive impairment, 1262
- Obstructive hypertrophic cardiomyopathy. *See* Cardiomyopathy
- Omega-3 fatty acids
- effects of dietary supplementation on endothelium-dependent vasodilation in chronic heart failure, 547
 - effects on men with healed MI and depressed ejection fractions
 - heart rate recovery after exercise, 1127
 - heart rate variability, 1127
 - resting heart rate, 1127
- Optical coherence tomography
- in assessment of coronary arterial plaques, 1172
 - in assessment of coronary thrombus, 1713
- Osteopenia, prevalence of atherosclerotic vascular disease in postmenopausal women with or without, 1427
- Osteoporosis, postmenopausal women with
- effects of raloxifene on cardiovascular adverse events in, 520
 - or without, prevalence of atherosclerotic vascular disease in, 1427
- Outflow tract reconstruction, vs Fontan repair, for pulmonary atresia/intact ventricular septum, exercise capacity after, 1638
- Outpatient echocardiography laboratory, impacts of obesity, 1082
- Overweight, patients with advanced heart failure, home-based exercise program for, 886
- Oxygen consumption, peak, vs ventilatory equivalent, in predicting outcomes of heart failure patients, 1607
- Pacing
- biventricular and left ventricular, effects on ventricular dyssynchrony in compensated heart failure, exercise echocardiographic assessment, 1622
 - dual-chamber, with echocardiographic evaluation of optimal atrioventricular delay in hypertrophic cardiomyopathy, 1769
 - long-term right ventricular, after His bundle ablation, left ventricular remodeling during, 1223
- new paradigm for cardiac rehabilitation resistance exercise guidelines, 281
- optimal left ventricular site, and tissue synchronization imaging in CRT, 1615
- Paclitaxel-eluting stents
- vs bare metal stents, in primary and rescue PCI for STEMI, 1308
 - vs sirolimus-eluting stents
 - in coronary arteries, late luminal loss as predictor of clinical efficacy, 603
 - for coronary in-stent restenosis, 1182
 - and dexamethasone-eluting stents, outcome, restenosis rates and CRP levels, 1311
 - effects on T-cell subset redistribution in CAD, 494
 - undeployed, failure to retrieve, 502
- PAI-1. *See* Plasminogen activator inhibitor 1
- Papillary fibroelastoma, of mitral valve, tumor excision vs valve replacement, 759
- Paroxetine, in depressed men with paroxysmal atrial fibrillation, 1749
- PAR-1 thrombin receptor. *See* Protease-activated receptor-1
- PBMV. *See* Percutaneous balloon mitral valvuloplasty
- PCI. *See* Percutaneous coronary intervention
- Pediatrics
- aortic root dilation and aortic regurgitation after tetralogy of Fallot repair, 905
 - children with heart failure due to idiopathic dilated cardiomyopathy
 - diastolic duration in, 101
 - systolic duration in, 101
 - MRI in diagnosis of arrhythmogenic right ventricular dysplasia, 1232
 - relation of carotid artery stiffness to obesity and metabolic syndrome, 528
- Percutaneous balloon mitral valvuloplasty, one-year outcome of cardioversion of atrial fibrillation in mitral valve stenosis after, 1045
- Percutaneous coronary intervention. *See also* Stents
- in ACS, intravascular ultrasound of coronary arteries during, 29
 - and adjunctive abciximab, ST segment analysis in AMI after, 48
 - cardiac catheterization laboratory and patient transfer for, 591 (Readers' Comments)
 - for chronic total coronary occlusions, trends in, 1691
 - coronary microvascular resistance during anterior wall AMI and after, 1131
 - drug-eluting vs bare metal stents for non-ST-elevation ACS, effects on biomarkers, 1473
 - effects of high vs low clopidogrel loading dose on stable angina pectoris patients subjected to, 984
 - elective high-risk, safety and feasibility of left ventricular support with Impella Recover LP 2.5, 990
 - first elective, effects of morning vs evening atorvastatin intake on major cardiac events and restenosis rates after, 44
 - high-risk, effects of GPRAs on inhibition of platelet aggregation in patients under, 489
 - hospital costs in treating complications of, 322
 - impact of ≥ 3 vs single sirolimus-eluting stents on outcomes in, 606
 - long-term aspirin and clopidogrel after, effects on platelet function, 38
 - MRI vs echocardiography to predict myocardial function improvement after, 361
 - multivessel vs single-vessel, outcomes of, 1585
 - neutrophil/lymphocyte ratio in prediction of long-term mortality after, 993
 - in patients with normal troponin T levels, N-Terminal ProBNP in prediction of long-term outcome after, 830
 - planned vs provisional use of GPRAs in smokers undergoing, 1679
- primary
- prediction of one-year mortality among 30-day STEMI survivors after, 817
 - and rescue, with paclitaxel-eluting stents vs bare metal stents for STEMI, 1308
- prognostic effects of distal vs proximal coronary occlusion on AMI after, 1137
- reactive hyperemia after, effect on myocardial tissue reperfusion in AMI, 617
- relation of CK-MB elevation and ischemic end points to angiographic adverse events during, 1591
- rescue
- after CABG in the drug-eluting stent era, 789
 - after failed thrombolysis, meta-analysis, 1685
- sirolimus-eluting stents vs bare metal stents
- impact of chronic renal insufficiency on outcomes of, 792
 - for saphenous vein graft, 34
- for STEMI, effect of platelet inhibition after abciximab administration on myocardial reperfusion after, 798

- stenting, for unstable angina, impact of clopidogrel on circulating levels of soluble CD40 ligand, 192
- vs thrombolysis, effects on outcomes of renal failure with STEMI, 1142
- usefulness and safety for cardiac transplant vasculopathy, 1192
- Percutaneous transluminal coronary angioplasty
- relation of ST-segment deviation to myocardial ischemia during acute coronary occlusion induced by, 295
 - vs stenting, cardiovascular complications of noncardiac surgery after, 1188
 - with vs without stenting, long-term outcomes, 1467
- Pericardial effusion, relation to fractional shortening, 910
- Peripartum cardiomyopathy. *See* Cardiomyopathy
- Peripheral arterial disease
- as marker of risk for abdominal aortic aneurysm in CAD, 1549 (Readers' Comments)
 - prevalence in postmenopausal women with or without osteoporosis or osteopenia, 1427
- Personality, type D, effects on prognosis of CHD, 970
- PET. *See* Positron emission tomography
- PISA. *See* Proximal isovelocity surface area
- Placebo effect
- on laser myocardial revascularization in CHD, 152 (Readers' Comments)
 - on therapeutic angiogenesis in CHD, 152 (Readers' Comments)
- Placenta-derived growth factors
- relation to development of collateral vessels, angiographic evaluation, 472
 - in stable angina pectoris and total coronary artery occlusion, 472
- Plant stanol, effects on low-density lipoprotein lowering in patients on statin therapy, 376
- Plaques
- coronary
 - assessment by optical coherence tomography, 1172
 - characterization using intravascular ultrasound vs multidetector computed tomography, 598
 - multiple unstable, manifested by recurrent STEMI in different territory after fibrinolysis, 947
 - relation to CHD, 1564
 - relation to left ventricular hypertrophy, effect of ethnicity, 1441
 - ruptured, in native coronary arteries vs saphenous vein grafts, intravascular ultrasound of, 593
- Plasminogen activator inhibitor 1, relation to depression and fatigue in patients with or without CHD, 1287
- Platelet aggregation
- inhibition after abciximab administration, effect on myocardial reperfusion after PCI for STEMI, 798
 - inhibition by GPRAs in patients undergoing high-risk PCI, 489
- Platelets
- in coronary thrombus of AMI patients, 26
 - expression of PAR-1 thrombin receptor, in patients with metabolic syndrome, effects of statins, 1332
 - function, effects of long-term aspirin and clopidogrel after PCI, 38
- Polymorphism
- of IGF-1 promoter, relation to mortality in elderly Caucasians with MI, 1274
 - of lipoprotein lipase, relation to dyslipidemia and diabetes in Asian Indians, 1337
- Positron emission tomography, fluorine-18 deoxyglucose, vs contrast-enhanced MRI, in detection of myocardial viability, 452
- Postmenopausal women. *See* Women
- Potassium, abnormal serum concentration, safety of exercise testing in patients with, 1247
- POTO. *See* Pulmonary outflow tract obstruction
- Pravastatin, benefits in CAD, effects of obesity, 1553
- Pregnancy, arrhythmia recurrence during, in women with previous tachyarrhythmia, impact on fetal and neonatal outcomes, 1206
- Pressure overload, secondary to congenital heart disease, effects on right ventricular function, 1051
- Primary care physician, vs disease management program, for idiopathic dilated cardiomyopathy, 253
- Prolapse, mitral valve, frequency of atrial fibrillation in mitral regurgitation secondary to, 1039
- Prosthesis failure
- assessment of leaflet motion in
 - transesophageal echocardiography in, 94
 - transthoracic echocardiography in, 94
 - clinical recurrence after sirolimus-eluting stent implantation, 1582
- Protease-activated receptor-1, platelet expression of, in patients with metabolic syndrome, effects of statins, 1332
- Proximal isovelocity surface area, vs transthoracic three-dimensional Doppler echocardiography, for assessment of mitral regurgitation, 1630
- Pseudoxanthoma elasticum, left ventricular systolic and diastolic functions in, 1535
- Psychological stress, effects on prognosis of CHD, 970
- PTCA. *See* Percutaneous transluminal coronary angioplasty
- Publications, good books in cardiology appearing in 2005, 430 (From the Editor)
- Pulmonary atresia, exercise capacity after univentricular vs biventricular repair for, 1638
- Pulmonary embolism
- acute
 - bedside testing for brain natriuretic peptides to identify right ventricular dysfunction and outcome in, 1386
 - predictors of hemorrhage after fibrinolysis for, 127
 - fatal, in ischemic stroke, 1776
- Pulmonary hypertension
- cardiopulmonary exercise testing in, 123
 - six-minute walk in, 123
- Pulmonary outflow tract obstruction, in neonates, significance of reverse orientation of ductus arteriosus for early intervention of, 716
- Pulmonary stenosis
- causing volume and/or pressure overload, effects on right ventricular function and B-type natriuretic peptides, 1051
 - peripheral, efficacy of medium-sized cutting balloons vs standard balloons for dilation of, 1060
- Pulmonary vein
- ablation, use of esophageal leads in determining strategies of, 1494
 - incorporation, relation to histology of myocardial wall of left atrium, 662
- P waves, during exercise testing, in CAD prediction, 1126 (Readers' Comments)
- QRS wave, in prediction of mortality after AMI, 353
- QT interval
- prolongation
 - by acute cocaine administration in habitual users, 1244
 - in ischemic or non-ischemic cardiomyopathy, effects of head-up tilt, 412
 - and subintervals, from large ethnically diverse population of women aged 50 to 79 years, 730
- Quality of health care
- in African-American patients admitted for CHF at a university teaching hospital, 690
 - patient and physician perceptions, 1073
- Quality of life, in CAD patients with or without diabetes mellitus, 1267
- Quinapril, effects on left atrial remodeling, left ventricular diastolic function and arterial stiffness, 916
- Q waves, redefinition of electrocardiographic criteria for MI, 974
- Racial differences. *See* Ethnicity
- Radiofrequency catheter ablation. *See* Catheter ablation
- Radionuclide imaging, myocardial ischemia during PTCA-induced acute coronary occlusion, relation to ST-segment deviation, 295
- Raloxifene, effect on cardiovascular adverse events in osteoporotic postmenopausal women, 520
- Real-time three-dimensional echocardiography. *See* Echocardiography
- Recovery of function, two weeks after AMI, 952
- Remodeling. *See also* Atrial remodeling; Ventricular remodeling
- patterns of minimally diseased coronary arteries, 13
 - postnatal, of ductus arteriosus, CD44-v6 expression in smooth muscle cells, 1056
- Renal failure, with STEMI, effects of PCI vs thrombolysis on outcomes of, 1142
- Renal impairment, gender differences in relation to angiographic CAD and long-term mortality, 630

- Renal insufficiency, chronic, impact on outcomes of PCI with sirolimus-eluting stents vs bare metal stents, 792
- Reperfusion
 in AMI, effect of reactive hyperemia after PCI on, 617
 myocardial
 after PCI for STEMI, effect of platelet inhibition after abciximab administration on, 798
 effect of pre-MI angina pectoris on arrhythmias after, 1157
 efficacy of, relation to N-Terminal ProBNP on admission of AMI patients, 1151
 thrombolysis vs PCI, effects on outcomes of renal failure with STEMI, 1142
- Restenosis
 after drug-eluting stent implantation
 vs brachytherapy, intravascular ultrasound of, 1292
 in one or more coronary arteries, predictors and incidence, 506
 after implantation of overlapping sirolimus-eluting (Cypher) stents, angiographic patterns, 499
 coronary in-stent, sirolimus vs paclitaxel-eluting stents for, 1182
 effects of morning vs evening atorvastatin intake after first elective PCI on, 44
 in-stent
 in diabetes mellitus patients, heterogeneity of neointimal distribution, 340
 on- vs off-label use of drug-eluting stents in clinical practice, 1478
 plasma aldosterone levels for prediction of, 785
 of saphenous vein graft, safety and efficacy of balloon angioplasty vs vascular brachytherapy vs drug-eluting stents for, 1303
 serial angiography after intracoronary brachytherapy for, 21
 rates after implantation of sirolimus- vs paclitaxel- vs dexamethasone-eluting stents, 1311
 in stable angina, soluble CD40 ligand in prediction of, 198
- Resynchronization. *See* Cardiac resynchronization therapy
- Retinal arteries, and CHD, 1549 (Readers' Comments)
- Revascularization
 in AMI, gender differences of, 1722
 effects of morning vs evening atorvastatin intake after first elective PCI on, 44
 incomplete, exercise myocardial scintigraphy in multivessel coronary disease after, 207
 laser myocardial, in CHD, placebo effect on, 152 (Readers' Comments)
 vs medical therapy, in CAD patients undergoing TAPSE, relation to survival, 804
 optimized, effects on incidence of sudden cardiac death after AMI, 480
 PTCA with vs without stenting, long-term outcomes, 1467
 of target vessel, in multivessel vs single-vessel PCI, 1585
- Rheumatic mitral stenosis, atrial fibrillation before isolated mitral valve replacement for, 1035
- Right atrium, pressure in anesthetized, paralyzed, and mechanically ventilated patients, accuracy of Doppler estimation, 1654
- Right ventricle
 in arrhythmogenic dysplasia
 in children, diagnosis using MRI, 1232
 feasibility and variability of MRI vs three-dimensional echocardiography, 703
 dysfunction
 in adult patients with operated tetralogy of Fallot, relation to elevated NT-pro-BNP and reduced exercise capacity, 1377
 and outcome in acute pulmonary embolism, bedside testing for brain natriuretic peptides for identification of, 1386
 function, effects of volume and/or pressure overload secondary to congenital heart disease, 1051
 involvement in Churg-Strauss syndrome, 1519
 long-term pacing, after His bundle ablation, left ventricular remodeling during, 1223
 peak systolic pressure, relation to major adverse events in CRT, 1737
 tissue Doppler imaging of free wall mechanical delay and dysfunction, 434 (Readers' Comments)
- Rosiglitazone, vs combination with atorvastatin, effects on biomarkers of cardiovascular disease in type 2 diabetes, 646
- Rosuvastatin
 vs atorvastatin, efficacy and safety in hypercholesterolemic African-Americans, 229
 short-term treatment, effect on estimated glomerular filtration rate, 1602
- Saphenous vein graft
 in-stent restenosis, safety and efficacy of balloon angioplasty vs drug-eluting stents, 1303
 vs vascular brachytherapy, 1303
 intravascular ultrasound of aorto-ostial vs shaft lesions, 1463
 vs native coronary arteries, intravascular ultrasound of ruptured plaques in, 593
 sirolimus-eluting vs bare metal stents for, 34
- Scintigraphy. *See* Radionuclide imaging
- Sclerosis, aortic valve, relation to morbidity and mortality in community-dwelling elderly individuals, 1281
- Self-transport, vs EMS, mortality in STEMI patients arriving by, 458
- Septal ablation, alcohol, for obstructive hypertrophic cardiomyopathy
 predictors of subacute complete heart block after, 264
 time course of pressure gradient response after, 1511
- Serotonin, coronary hypersensitivity to, in cardiac transplant recipients, effects of vitamin C, 561
- Sibutramine, short-term therapy, relation to weight loss and improved endothelial function in obese patients with CAD, 1650
- Simvastatin
 monotherapy, vs Vytorin, muscle safety profiles, 223
 phenotypic predictors of response to, African-Americans vs Caucasians, 843
- Single-photon emission computed tomography
 gated, relation of postexercise abnormal heart rate recovery to left ventricular perfusion and function, 1452
 infarct size, relation to N-Terminal ProBNP on admission of AMI patients, 1151
 for MI and/or coronary revascularization, 1
 vs myocardial contrast echocardiography, to predict AMI transmural, 1718
 stress technetium-99m-tetrofosmin myocardial perfusion, 1
 for cardiovascular disease risk stratification in metabolic syndrome and diabetes, 1538
 in multivessel coronary disease after incomplete revascularization, 207
 technetium-99m sestamibi, effect of body mass index on left ventricular cavity size and ejection fraction, 725
- Sirolimus-eluting stents
 vs bare metal stents
 impact of chronic renal insufficiency on outcomes of PCI with, 792
 safety and efficacy for STEMI, 1299
 for saphenous vein graft intervention, 34
 characterization of the "black hole" phenomenon after, 203
 for coronary bifurcation lesions, implantation by crush technique, long-term outcomes, 1317
 overlapping, angiographic patterns of restenosis after implantation of, 499
 vs paclitaxel-eluting stents
 in coronary arteries, late luminal loss as predictor of clinical efficacy, 603
 for coronary in-stent restenosis, 1182
 vs dexamethasone-eluting stents, outcome, restenosis rates and CRP levels, 1311
 effects on T-cell subset redistribution in CAD, 494
 safety and feasibility of various implantation techniques for unprotected left main coronary artery bifurcation stenosis, 1597
 ≥ 3 vs single, impact on outcomes in PCI patients, 606
- Six-minute walk
 and cardiopulmonary exercise testing, 123
 in pulmonary hypertension, 123
- Smokers, undergoing pCI, planned vs provisional use of GPRAs in, 1679
- Smooth muscle cells, in postnatal remodeling of ductus arteriosus, CD44-v6 expression in, 1056
- Social integration, relation to CRP and IL-6 levels in men and women 70 to 79 years, 1010
- Sodium, serum levels, effects of tolvaptan vs fluid restriction, 1064
- SPECT. *See* Single-photon emission computed tomography
- Sphygmograph, in America, historical study, 580
- Stable angina. *See* Angina pectoris
- Stanol, plant, effects on low-density lipoprotein lowering in patients on statin therapy, 376
- Statins
 adding colesvelam hydrochloride to, in hypercholesterolemia
 effects on high-sensitivity CRP, 1198

- effects on low-density lipoprotein, 1198
- in cholesterol management by targeting lower low-density lipoprotein concentration, 1667 (Editorial)
- effect on coronary collateral development in diabetes mellitus, 772
- effect on infarct-related coronary artery patency, 7
- effect on platelet expression of PAR-1 thrombin receptor in patients with metabolic syndrome, 1332
- effect on risk
 - of atrial fibrillation after CABG, 55, 1266 (Readers' Comments), 1551 (Readers' Comments)
 - of ventricular arrhythmia in CAD, 153 (Readers' Comments)
- effects of plant stanol on low-density lipoprotein lowering in patients on, 376
- effects on CHF and myocardial ischemia in Japanese AMI patients, 1165
- effects on kidney function, 748
- effects on mortality and size of abdominal aortic aneurysm, 279
- effects on renal function after aortic cross clamping during major vascular surgery, 1383
- influence on length of hospitalization in major vascular surgery patients, 1423
- over-the-counter availability, impact on CHD, 851
- time of initiation after hospital discharge
 - relation to mortality in AMI patients ≥ 65 years of age, 155
 - relation to risk of recurrent AMI in patients ≥ 65 years of age, 155
- use before STEMI, 7
- STEMI. *See* ST-segment elevation myocardial infarction
- Stenosis. *See also* Coronary stenosis
 - aortic valve
 - CRP as predictor of progression of, 90
 - levels of aldosterone, angiotensin II and B-type natriuretic peptide in ascending aorta and coronary sinus, 1068, 1107 (Editorial)
 - mitral valve, after PBMV, one-year outcome of cardioversion of atrial fibrillation in, 1045
 - peripheral pulmonary, efficacy of medium-sized cutting balloons vs standard balloons for dilation of, 1060
 - pulmonary, causing volume and/or pressure overload,
 - effects on B-type natriuretic peptides, 1051
 - effects on right ventricular function, 1051
 - rheumatic mitral, atrial fibrillation before isolated mitral valve replacement for, 1035
 - unprotected left main coronary artery bifurcation, safety and feasibility of various sirolimus-eluting stent implantation techniques for, 1597
- Stents
 - vs balloon angioplasty, outcomes of AMI after, 977
 - bare metal, circadian variation in symptomatic subacute stent thrombosis, 195
 - carotid artery, optimal activated clotting time during implantation of, 1657
 - "direct," without balloon predilatation, effectiveness of, 316
 - drug-eluting
 - vs bare metal, for non-ST-elevation ACS, effects on biomarkers, 1473
 - vs brachytherapy, intravascular ultrasound of restenosis after, 1292
 - on- vs off-label use in clinical practice, 1478
 - for ostial de novo left anterior descending coronary artery lesions, early outcome, 187
 - predictors and incidence of restenosis after placement in one or more coronary arteries, 506
 - rescue PCI after CABG, 789
 - vs vascular brachytherapy vs balloon angioplasty, safety and efficacy for saphenous vein graft in-stent restenosis, 1303
 - exercise myocardial scintigraphy in multivessel coronary disease after incomplete revascularization with, 207
 - implantation in AMI, angiographic no-reflow phenomenon during, 301
 - long-term outcomes of PTCA with vs without, 1467
 - paclitaxel-eluting. *See* Paclitaxel-eluting stents
 - in patients with unstable angina pectoris, impact of clopidogrel on circulating levels of soluble CD40 ligand, 192
 - vs PTCA, cardiovascular complications of noncardiac surgery after, 1188
 - restenosis
 - in diabetes mellitus patients, heterogeneity of neointimal distribution in, 340
 - plasma aldosterone levels for prediction of, 785
 - serial angiography after intracoronary brachytherapy for in-stent restenosis, 21
 - sirolimus-eluting. *See* Sirolimus-eluting stents
 - for stable angina, soluble CD40 ligand in prediction of restenosis, 198
 - Strain, two-dimensional and tissue based, validation with harmonic phase MRI, 1661
 - Stress
 - blood pressure reactivity to, and left ventricular mass among Caucasian and Africa-American men and women, 240
 - effects on prognosis of CHD, 970
 - intramyocardial injection of autologous bone marrow-derived mononuclear cells for myocardial ischemia induced by, 1326
 - Stress testing
 - after CABG, 810
 - dobutamine echocardiography
 - heart rate reserve vs 85% of age-predicted maximum heart rate as measure of chronotropic response during, 742
 - in observing wall motion abnormalities during recovery after acute beta-blocker administration, 462
 - for MI and/or coronary revascularization, 1
 - myocardial perfusion
 - vs coronary calcium, in apparently healthy siblings of individuals with premature CAD, 328
 - relation of myocardial ischemia to ventricular premature complexes during recovery from, 1570
 - safety in patients with abnormal serum potassium concentration, 1247
 - technetium-99m-tetrofosmin myocardial perfusion, 1
 - for cardiovascular disease risk stratification in metabolic syndrome and diabetes, 1538
 - Stroke
 - ischemic, fatal pulmonary embolism in, 1776
 - prediction by aortic root dimension in persons ≥ 65 years of age, 270
 - prevalence in postmenopausal women with or without osteoporosis or osteopenia, 1427
 - ST-segment analysis, in AMI after PCI and adjunctive abciximab, 48
 - ST-segment deviation
 - in lead aVR, and troponin T, for early risk stratification in non-ST-segment elevation ACS, 334
 - relation to myocardial ischemia during PTCA-induced acute coronary occlusion, 295
 - ST-segment elevation, for predicting outcomes after primary coronary angioplasty vs fibrinolysis, 611
 - ST-segment elevation myocardial infarction
 - 30-day survivors after primary PCI, prediction of one-year mortality among, 817
 - effect of platelet inhibition after abciximab administration on myocardial reperfusion after PCI for, 798
 - effects of PCI vs thrombolysis on outcomes of renal failure with, 1142
 - medication use prior to, effect on infarct-related coronary artery patency, 7
 - mortality in patients arriving by EMS vs self-transport, 458
 - nocturnal melatonin levels and CRP in patients with, 10
 - on- vs off-label use of drug-eluting stents in clinical practice, 1478
 - paclitaxel-eluting stents vs bare metal stents in primary and rescue PCI for, 1308
 - prognostic value of nocturnal melatonin levels, 1162
 - recurrence in different territory, after fibrinolysis, as manifestation of multiple unstable coronary arterial plaques, 947
 - safety and efficacy of sirolimus-eluting stents vs bare metal stents for, 1299
 - Sudden death
 - incidence after AMI, effects of optimized medical and revascularization therapy, 480
 - in young female military recruits, 1756
 - Surgery
 - cardiac, incidence and risk factors of heart failure in adolescents and adults with congenital heart disease after, 1238
 - clinician use of perioperative cardiovascular testing guidelines, effects on outcomes, 118
 - major vascular
 - effects of statins on renal function after aortic cross clamping during, 1383
 - influence of cardiac risk factors and medication length of hospitalization, 1423

- noncardiac
 after PTCA vs stenting, cardiovascular complications of, 1188
 prognostic value of routine preoperative electrocardiography in, 1103
 valve, detection of coronary stenosis with multidetector computed tomography vs noninvasive coronary angiography before, 1506
- Survival
 after AMI, effect of β -blockers, 1125 (Readers' Comments)
 after MI among diabetics, racial differences on, 1551 (Readers' Comments)
 long-term, relation to revascularization vs medical therapy in CAD patients undergoing TAPSE, 804
 in patients with normal left ventricular ejection fractions, 959
 Swan, Harold James Charles, 1545 (Historical Study)
 Sympathy, and empathy, clinical usefulness of, 1789 (Editorial)
- Syncope
 high risk of ventricular arrhythmias in patients with nonischemic dilated cardiomyopathy with, 416
 neurally mediated, contribution of decreased atrial function in pathogenesis of, 1017
- Systolic function
 in children with heart failure due to idiopathic dilated cardiomyopathy, 101
 vs diastolic function
 in heart failure patients ≥ 65 years of age, left atrium in, 83
 left ventricular, as outcome predictor after PCI for AMI, 160
 in heart failure, prognostic value of restrictive mitral filling pattern, 676
 left ventricular
 after exercise training in young men, 1089
 in Behçet's disease, pulse-wave Doppler and Doppler tissue imaging of, 716
 effects of growth hormone therapy, 1097
 effects of long-term cocaine use, 1085
 effects of tuna or other broiled or baked fish vs fried fish, 216
 impaired, prediction of exercise tolerance by diastolic mitral annular velocity, 1025
 in ischemic or idiopathic dilated cardiomyopathy, tissue Doppler assessment of ejection fractions and IVA, 872
 in non-compacted and compacted segments in isolated left ventricular non-compaction, 738
 in pseudoxanthoma elasticum, 1535
 relation to pericardial effusion, 910
 remodeling in organic and functional mitral regurgitation, relation to B-type natriuretic peptide activation, 1029
 peak pressure, right ventricular, relation to major adverse events in CRT, 1737
- Tachyarrhythmia. *See* Tachycardia
- Tachycardia, previous, arrhythmia recurrence during pregnancy in women with, impact on fetal and neonatal outcomes, 1206
- Tadalafil, for erectile dysfunction, incidence of cardiovascular adverse events, 1778
- TAPSE. *See* Transesophageal atrial pacing stress echocardiography
- Taxus stents, undeployed, failure to retrieve, 502
- Technetium-99m sestamibi, SPECT, effect of body mass index on left ventricular cavity size and ejection fraction, 725
- Technetium-99m-tetrofosmin
 exercise scintigraphy, in multivessel coronary disease after incomplete revascularization, 207
 SPECT, in MI and/or coronary revascularization, 1
 stress SPECT, for cardiovascular disease risk stratification in metabolic syndrome and diabetes, 1538
- Tei index, relation of biventricular dysfunction to elevated NT-pro-BNP and reduced exercise capacity in adult patients with operated tetralogy of Fallot, 1377
- Tetralogy of Fallot
 causing volume and/or pressure overload, effects on right ventricular function and B-type natriuretic peptides, 1051
 operated, relation of biventricular dysfunction to elevated NT-pro-BNP and reduced exercise capacity in adult patients with, 1377
 repair, aortic root dilation and aortic regurgitation after, 905
- TF. *See* Tissue factor
- Thiazolidinediones, effect on serum CRP levels, meta-analysis, 655
- Thomas, Vivien, 435 (Readers' Comments)
- Three-dimensional echocardiography. *See* Echocardiography
- Thrombocytopenia, prevalence and management with GPRAs, 426
- Thromboembolism, left atrial appendage, detection by transesophageal echocardiography 421
- Thrombolysis
 effects during out-of-hospital CPR, 305
 failed, meta-analysis of rescue PCI after, 1685
 vs PCI, effects on outcomes of renal failure with STEMI, 1142
 Thrombolysis In Myocardial Infarction risk score, during ER stay in low-risk patients with acute chest pain, frequency and timing of changes in, 781
- Thrombosis, symptomatic subacute stent, after bare metal coronary stent implantation, circadian variation in, 195
- Thrombotic endocarditis, nonbacterial, in cancer and cerebral ischemia, detection by transesophageal echocardiography, 894
- Thyroid gland, intracardiac accessory, 926
- TIMI risk score. *See* Thrombolysis In Myocardial Infarction risk score
- TIMPs. *See* Tissue inhibitors of metalloproteinases
- Tirofiban
 effects on inhibition of platelet aggregation in patients undergoing high-risk PCI, 489
 for thrombocytopenia, 426
- Tissue factor, in coronary thrombus of AMI patients, 26
- Tissue inhibitors of metalloproteinases, selective induction in atrial and ventricular myocardium in patients with atrial fibrillation, 532
- Tissue plasminogen activator antigen, relation to depression and fatigue in patients with or without CHD, 1287
- Tissue synchronization imaging, and optimal left ventricular pacing site in CRT, 1615
- T-lymphocytes, subset redistribution in CAD, effects of sirolimus- vs paclitaxel-eluting stents, 494
- Tolvaptan, vs fluid restriction, for hyponatremia, 1064
- t-PA antigen. *See* Tissue plasminogen activator antigen
- Transesophageal atrial pacing stress echocardiography, relation of survival to revascularization vs medical therapy in CAD patients undergoing, 804
- Transesophageal echocardiography
 in assessment of leaflet motion in prosthetic valve dysfunction, 94
 in patients with cancer and cerebral ischemia, for detection of cardiac sources of embolism, 894
 of nonbacterial thrombotic endocarditis, 894
 in prediction of left atrial appendage thrombus 421
 tissue Doppler, for assessment of aortic wall mechanics in Marfan syndrome, 571
 vs transthoracic, diagnosis of right-to-left atrial shunting, 434 (Readers' Comments)
- Transthoracic echocardiography
 in assessment of leaflet motion in prosthetic valve dysfunction, 94
 three-dimensional Doppler, vs PISA, for assessment of mitral regurgitation, 1630
 vs transesophageal, diagnosis of right-to-left atrial shunting, 434 (Readers' Comments)
- Triglyceride/high-density lipoprotein ratio, in prediction of low-density lipoprotein particle size, phenotype B and particle concentrations in Asian Indians, 1007
- Troponin, chest pain positive for, contrast-enhanced MRI in detection of myocardial scar in, 768
- Troponin T
 normal levels, N-Terminal ProBNP in prediction of long-term outcome after PCI in patients with, 830
 for predicting long-term mortality
 in diabetes mellitus, 167
 in non-ST-elevation ACS, 167
 and ST segment in lead aVR, for early risk stratification in non-ST-segment elevation ACS, 334
- Tumor excision, vs valve replacement, for papillary fibroelastoma of mitral valve, 759
- Turner syndrome, effects of growth hormone treatment on aortic distensibility and dimensions in, 1644
- T waves
 abnormalities, relation to extent of CAD, 1124 (Readers' Comments)

- alternans, relation to regional left ventricular dysfunction and eccentric hypertrophy secondary to CHD, 775
in prediction of mortality after AMI, 353
- Twins, identical, with hypertrophic cardiomyopathy and left ventricular apical aneurysms, 1109
- Type 1 diabetes mellitus. *See* Diabetes mellitus, type 1
- Type 2 diabetes mellitus. *See* Diabetes mellitus, type 2
- Type D personality, effects on prognosis of CHD, 970
- Ultrasound, intravascular. *See* Intravascular ultrasound
- Unstable angina. *See* Angina pectoris
- Valve prosthesis
adherence to ACC/AHA guidelines on periprocedural anticoagulation in patients with, 891
in assessment of leaflet motion in dysfunction of
transesophageal echocardiography, 94
transthoracic echocardiography, 94
influence on erythrocyte survival, assessment by breath carbon monoxide, 1374
mitral, Starr-Edwards model 6120, 756
vs tumor excision, for papillary fibroelastoma of mitral valve, 759
warfarin-related over-anticoagulation in patients with
FFP for, 409
vitamin K for, 409
- Valvuloplasty, percutaneous balloon mitral, one-year outcome of cardioversion of atrial fibrillation in mitral valve stenosis after, 1045
- Vascular adhesion molecule-1, soluble, serum levels in patients with coronary vasospasm without CAD vs patients with stable angina and ACS with CAD, 1429
- Vascular endothelial growth factor
relation to development of collateral vessels, angiographic evaluation, 472
in stable angina pectoris and total coronary artery occlusion, 472
- Vasculopathy. *See* Coronary artery disease
- Vasodilation
endothelium-dependent, in chronic heart failure, effects of dietary omega-3 fatty acid supplementation on, 547
metabolic, in response to exercise, ischemia and endothelium-dependent flow-mediated dilation in African-American vs non-African-American patients with chronic heart failure, 685
- Vasospasm, coronary
induced by ergonovine maleate in vasospastic angina pectoris, intravascular ultrasound of, 1322
without CAD, vs stable angina and ACS with CAD, serum levels of inflammatory markers in patients with, 1429
- Vegetarian diet. *See* Diet
- Ventilatory equivalent, vs peak oxygen exercise consumption, in predicting outcomes of heart failure patients, 1607
- Ventricle
myocardium in patients with atrial fibrillation, selective induction of TIMPs and MMP in, 532
volume and function, in persistent atrial fibrillation before and after cardioversion, 1213
- Ventricular arrhythmias. *See* Arrhythmias
- Ventricular dysfunction
left
in adult patients with operated tetralogy of Fallot, relation to elevated NT-pro-BNP and reduced exercise capacity, 1377
secondary to CHD, relation to T-wave alternans, 775
right
in adult patients with operated tetralogy of Fallot, relation to elevated NT-pro-BNP and reduced exercise capacity, 1377
and outcome in acute pulmonary embolism, bedside testing for brain natriuretic peptides for identification of, 1386
- Ventricular dyssynchrony, in compensated heart failure, exercise echocardiographic assessment of effects of biventricular and left ventricular pacing, 1622
- Ventricular fibrillation, in users of implantable cardioverter-defibrillator, prevalence and predictors of, 389
- Ventricular function. *See* Left ventricular function; Right ventricular function
- Ventricular premature complexes, during recovery from myocardial perfusion exercise stress test, relation to myocardial ischemia, 1570
- Ventricular remodeling
end-systolic left, in organic and functional mitral regurgitation, relation to B-type natriuretic peptide activation, 1029
left
in CHF patients, effect of CRT on reversal of, 876
in chronic severe mitral regurgitation, plasma N-terminal ProBNP as marker of, 1498
during long-term right ventricular pacing after His bundle ablation, 1223
- Ventricular tachycardia, in users of implantable cardioverter-defibrillator, prevalence and predictors of, 389
- Vital exhaustion. *See* Fatigue
- Vitamin C, effects on coronary hypersensitivity to serotonin in cardiac transplant recipients, 561
- Vitamin K, efficacy and safety for warfarin-related over-anticoagulation in patients with mechanical heart valves, 409
- Volume overload, secondary to congenital heart disease, effects on right ventricular function, 1051
- von Willebrand factor, in coronary thrombus of AMI patients, 26
- Vytorin, vs simvastatin monotherapy, muscle safety profiles, 223
- Wait-to-hip ratio, impact on systemic hypertension in lean Chinese, 839
- Warfarin
anticoagulation, effectiveness of manual pressure hemostasis after transfemoral coronary angiography in patients on, 485
over-anticoagulation, in patients with mechanical heart valves, efficacy and safety of vitamin K and FFP for, 409
use in patients with new-onset atrial fibrillation, patterns and predictors of, 538
- Weight loss, in obese patients with CAD, relation to short-term sibutramine therapy, 1650
- White blood cell count
and body mass index, relation to metabolic syndrome, 835
for predicting in-hospital outcomes after AMI, 1558
- Women
aged 50 to 79 years, QT interval and subintervals from ethnically diverse population of, 730
all-cause mortality risk vs CHD for electrocardiographic abnormalities in, 309
with cardiac syndrome X, microvascular function, metabolic syndrome and risk factor status in, 1727
Caucasian and Africa-American, blood pressure stress reactivity and left ventricular mass among, 240
healthy postmenopausal, effects of vegetarian diet on
blood pressure in, 380
HRV in, 380
and men, 70 to 79 years, relation of social integration to CRP and IL-6 levels, 1010
osteoporotic postmenopausal, effects of raloxifene on cardiovascular adverse events in, 520
postmenopausal
with or without osteoporosis or osteopenia, prevalence of atherosclerotic vascular disease in, 1427
prevalence and determinants of electrocardiographic left ventricular hypertrophy in multiethnic population of, 512
with previous tachyarrhythmia, arrhythmia recurrence during pregnancy in, impact on fetal and neonatal outcomes, 1206
with suspected CAD, estrogen status as prognostic marker in, 367
young military recruits, sudden death in, 1756
- Women, Hispanic
impaired endothelial function among, 68
increased body mass index among, 68
- Work, return to, two weeks after AMI, 952
- Yawning, repetitive, in cardiac tamponade, 1547 (Readers' Comments)