

- Academic Research Consortium, incidence of drug-eluting stent thrombosis based on definitions of, 683
- ACE inhibitors. *See* Angiotensin-converting enzyme inhibitors
- Acute aortic dissection
 risk factors for diagnostic delay in, 1399
 type A, geographic differences in clinical presentation, treatment, and outcomes in (from the International Registry of Acute Aortic Dissection), 1562
- Acute coronary events. *See also* Cardiac events
 Global Registry of Acute Coronary Events (GRACE) risk score on, 1114 (Readers' Comments)
 vulnerable plaques and, 1644 (Editor's Roundtable)
- Acute coronary syndrome
 admission neutrophil to lymphocyte ratio and outcomes in patients with, 653
 distribution and frequency of thin-capped fibroatheromas and ruptured plaques in the entire culprit coronary artery as determined by optical coherence tomography in patients with, 975
 in hospitalized patients, prognostic value of free plasma homocysteine levels with, 135
 impact of optimal medical therapy and revascularization on outcome of patients with chronic kidney disease and on dialysis with, 535
 magnitude and prognosis of ventricular arrhythmias associated in patients hospitalized with (from the GRACE Registry), 1577
 non-ST-segment elevation
 intensive atorvastatin therapy effect on prostaglandin E2 levels and metalloproteinase-9 activity in the plasma of patients with, 12
 outcomes of patients with previous coronary artery bypass grafting and (from the Pravastatin or Atorvastatin Evaluation and Infection Therapy [PROVE IT-TIMI 22] and the Aggrastat to Zocor [A to Z] Trials), 552
 percutaneous coronary interventions for, clopidogrel benefit doubt due to rebound adverse events after, 248 (Readers' Comments)
 percutaneous coronary intervention versus coronary artery bypass grafting as primary revascularization in patients with, 173
 pregnancy associated plasma protein A as a quick and sensitive biomarker in early phase of, 954 (Readers' Comments)
 prognostic implications of normal (<0.10 ng/ml) and borderline (0.10 to 1.49 ng/ml) troponin elevation levels in critically ill patients without, 509
 weekend/holiday versus weekday hospital discharge and guideline adherence on (from the American Heart Association's Get With the Guidelines – Coronary Artery Disease Database), 663
- Acute Decompensated Heart Failure National Registry Emergency Module, illicit stimulant use in a United States heart failure population presenting to the emergency department, 1216
- Acute myocardial infarction. *See also* ST-segment elevation myocardial infarction
 bans on smoking in public places and incidence of, 1421 (Editorial)
 blood transfusion and mortality risk in, 1116 (Readers' Comments)
 chest pain center accreditation is associated with better performance of Centers for Medicare and Medicaid Services core measures for, 120
 comparison of value of leads from body surface maps to 12-lead electrocardiogram for diagnosis of, 257
 complicated by ventricular fibrillation, thirty-year trends (1975–2005) in the magnitude, patient characteristics, and hospital outcomes of patients with, 1595
 failure to prescribe beta blockers acutely in patients with sustained ventricular tachycardia and ventricular fibrillation following, mortality and (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry), 1427
 invasive coronary revascularization effect on subsequent death rate and frequency of chronic heart failure in, 1
 left ventricular conic index measured by real-time three-dimensional echocardiography to predict left ventricular remodeling after, 1433
 of left ventricular wall, right ventricular cardiomyocyte apoptosis in patients with, 658
 posterior thorax data from body surface maps and standard electrocardiograms in patients with, 1117 (Readers' Comments)
 red blood cell transfusion impact on clinical outcomes in patients with, 115
 relation between previous lipid-lowering therapy and infarct size (creatinine kinase-MB level) in patients presenting with, 1119
 reloading effectiveness to overcome clopidogrel nonresponsiveness in patients with, 524
 seasonal pattern of incidence and case fatality in a Japanese population with (from the Takashima AMI Registry, 1988 to 2003), 1307
 socioeconomic status influence on lifestyle modifications among survivors of, 1583
 statin therapy and risk of heart failure after, 1706
 trends in prehospital delay in patients with (from the Worcester Heart Attack Study), 1589
- Adenosine, heart rate response in patients with diabetes mellitus to, normal myocardial perfusion imaging and, 1103
- Adhesion molecules, in hypertensive and normotensive postmenopausal women, effect of soy nuts on inflammation markers and, 84
- Adipose tissue, epicardial, coronary atherosclerosis and, 1602
- Adolescents
 autonomic heart rate control at rest and during unloading of the right ventricle in repaired tetralogy of Fallot in, 1085
 with hypertrophic cardiomyopathy, significance of left atrial volume in, 1390
- Adults. *See also* Aging; Elderly
 with complex congenital heart disease and ventricular dysfunction (subaortic right ventricle and single ventricle physiology), prevalence and determinants of, 625
 healthy young
 carnitine supplementation effect on flow-mediated dilation and vascular inflammatory responses to a high-fat meal in, 1413
 sleep apnea, atrial fibrillation, and heart failure implications from dynamic changes of left ventricular performance and left atrial volume induced using the Mueller maneuver in, 1557
 incidence of atrial fibrillation following transcatheter closure of atrial septal defects in, 902
 serum cystatin C and increased prevalence of coronary heart disease without chronic kidney disease in, 54
- Aerobic exercise testing, in male firefighters, comparison of cardiocirculatory and thermal strain during fire suppression to, 1551
- Aerobic interval-training program, group-based high-intensity, in patients with chronic heart failure, long-term effects of, 1220, 1361
- African Americans. *See also* Blacks
 epidemiology, heritability, and genetic linkage of C-reactive protein in (from the Jackson Heart Study), 835
- Age factors
 double product (systolic blood pressure × heart rate) reserve-adjusted modification of the Duke Treadmill Score nomogram in men and, 1407
 post cardiac catheterization contrast-induced acute kidney injury, gender and, 1482
- Aggrastat to Zocor (A to Z) Trials, outcomes of patients with previous coronary artery bypass grafting, 552
- Aging
 adverse effects on carotid artery intima-media thickness of, parental coronary artery disease and (from the Bogalusa Heart Study), 180
 caffeinated coffee consumption, cardiovascular disease, and heart valve disease in (from the Framingham Heart Study), 1502
 chronic kidney disease, diabetes mellitus, and myocardial infarction as equivalent risk factors for cardiovascular mortality with, 1668
- Air pollution, fine particulate, heart failure hospitalization and exposure to, 1230
- Alcohol
 consumption, and heart failure in hypertensive US male physicians, 593
 low-doses of, and warm-up phenomenon in patients with stable angina pectoris, 146
- Alcohol septal ablation, for hypertrophic cardiomyopathy, and electrocardiographic pattern in, 621
- Aldosterone synthase, gene polymorphism as atrial fibrillation determinant in patients with heart failure, 326

- Ambulatory patients, with coronary heart disease, left atrial volume index versus left ventricular ejection fraction usefulness to predict heart failure hospitalization and mortality in (from the Heart and Soul Study), 70
- American College of Cardiology/American Heart Association practice guidelines on bicuspid aortic valve with dilated ascending aorta, critical review of, 107
- practice guidelines on bicuspid aortic valve with dilated ascending aorta, critical review of the critical review of, 111 (Editorial)
- American Heart Association's Get With the Guidelines – Coronary Artery Disease Database
- weekend/holiday versus weekday hospital discharge and guideline adherence with acute coronary syndrome, 663
- American Journal of Cardiology*, proceedings of the Editorial Board Meeting on 30 March 2008, 231 (From the Editor)
- AMI. *See* Acute myocardial infarction
- Anemia
- in adults with complex congenital heart disease and ventricular dysfunction (subaortic right ventricle and single ventricle physiology), prevalence and determinants of, 625
- comparison of blood volume characteristics in patients with low versus preserved left ventricular ejection fractions and, 1069
- as predictor for presence and severity of coronary artery disease in patients undergoing stress myocardial perfusion study, effectiveness of, 266
- Aneurysmal subarachnoid hemorrhage, relation of cardiac troponin I to clinical severity, cardiac dysfunction, and pulmonary congestion in patients with, 1545
- Angina pectoris
- prognostic significance of myocardial ischemia by dobutamine stress echocardiography after coronary revascularization in patients without, 1156
- in a referral population, profile of chronic and recurrent, 1301
- stable, low doses of alcohol and warm-up phenomenon in patients with, 146
- stable or unstable, bleeding risk and outcomes of bivalirudin versus glycoprotein IIb/IIIa inhibitors with targeted low-dose unfractionated heparin in patients having percutaneous coronary intervention for, 160
- unstable, hemorheological variables in ST-elevation myocardial infarction or non-ST-elevation myocardial infarction versus, 125
- Angiography. *See* Coronary angiography
- Angioplasty. *See* Coronary angioplasty
- Angiotensin-converting enzyme (ACE) inhibitors
- digoxin and clinical outcomes in systolic heart failure patients on, 1356
- mortality in patients with implantable cardioverter-defibrillators and, 77
- Angiotensin receptor blockers
- cardiovascular protection and, from the Ongoing Telmisartan Alone and in Combination With Ramipril Global Endpoint Trial (ONTARGET), 1281 (Editorial)
- mortality in patients with implantable cardioverter-defibrillators and, 77
- Anticoagulant therapy
- long-term oral, perioperative bridging therapy with unfractionated heparin or low-molecular-weight heparin in mechanical prosthetic valve patients on (from the REGIMEN Registry), 879
- long-term outcomes in patients undergoing coronary stenting on dual oral antiplatelet treatment requiring, 1618
- Antiplatelet therapy
- dual oral, requiring oral anticoagulant therapy, long-term outcomes in patients undergoing coronary stenting on, 1618
- prolonged dual (aspirin + thienopyridine), incidence of bleeding and compliance following drug-eluting stent implantation with, 1477
- upstream patterns before primary percutaneous coronary intervention for acute ST-elevation myocardial infarction of (from the CRUSADE National Quality Improvement Initiative), 1335
- Aorta
- abdominal, percutaneous, minimally invasive retrieval of dislodged and impacted atrial septal defect closure device from, 1111
- calcium level, Framingham risk score for calcium level in coronary arteries or carotid artery intima-media thickness versus, 825
- descending, continuity equations in increased Doppler velocities due to aortic regurgitation in, 372 (Readers' Comments), *Reply*, 372
- dilated, absence of TGFBR1 and TGFBR2 mutations in patients with bicuspid aortic valve and, 629
- dilated ascending
- with bicuspid aortic valve
- critical review of American College of Cardiology/American Heart Association practice guidelines for, 107
- critical review of the critical review of the American College of Cardiology/American Heart Association practice guidelines for, 111 (Editorial)
- distensibility in patients with bicuspid aortic valves, 370 (Readers' Comments)
- Aortic regurgitation
- with central versus eccentric jets, flow convergence method accuracy for quantification in patients with, 475
- continuity equations in increased Doppler velocities in the descending aorta due to, 372 (Readers' Comments), *Reply*, 372
- Aortic valve(s)
- bicuspid
- absence of TGFBR1 and TGFBR2 mutations in patients with aortic dilation and, 629
- aortic distensibility in patients with, 370 (Readers' Comments)
- with dilated ascending aorta
- critical review of American College of Cardiology/American Heart Association practice guidelines for, 107
- critical review of the critical review of the American College of Cardiology/American Heart Association practice guidelines for, 111 (Editorial)
- bileaflet mechanical prostheses or bioprostheses, Doppler–catheter discrepancies in patients with, 1383
- isolated replacement for aortic stenosis, usefulness of left ventricular shape to predict early recovery of left ventricular function after, 1530
- percutaneous implantation of, feasibility and initial results including selection of transfemoral or transapical approach in patients with severe aortic stenosis, 1240
- replacement for aortic stenosis, mitral regurgitation and, 1378
- replacement for symptomatic aortic stenosis, comparison of brain natriuretic peptide plasma levels versus logistic EuroSCORE in predicting in-hospital and late postoperative mortality in patients undergoing, 749
- stage-related effect of statin treatment on the progression of sclerosis and stenosis of, 738
- stenosis, asymptomatic, atorvastatin modulation of prognosis and risk factors in patients with, 743
- Aortic wall, thickness and distensibility relation to cardiovascular risk factors (from the Multi-Ethnic Study of Atherosclerosis [MESA]), 491
- Arrhythmias
- diagnosis of accessory pathway using the His-His (H-H) and atrial-atrial (A-A) intervals of the first entrained cycle during ventricular overdrive pacing for, 197
- impact of β 1-adrenergic receptor polymorphisms on prognosis, response to beta-blocker therapy, susceptibility to heart failure and, 726
- prediction in ischemic cardiomyopathy beyond one year using microvolt T-wave alternans, 280
- Aspirin
- dose of, in patients with recent cerebrovascular or coronary ischemic events, relation between bleeding, all-cause mortality, and (from the BRAVO Trial), 1285
- low-dose, comparison of increased aspirin dose versus combined aspirin plus clopidogrel therapy in patients with diabetes mellitus and coronary heart disease and impaired antiplatelet response to, 396
- and thienopyridine following drug-eluting stent implantation, incidence of bleeding and compliance with, 1477
- Asthma
- adult-onset, relation to coronary heart disease and stroke, 508 (Readers' Comments)
- cardiac event risk in patients with long-QT syndrome treated with β_2 agonists and, 866
- Atherosclerosis, coronary
- detected by multislice computed tomography, association of traditional risk-factor assessment with, 316
- epicardial adipose tissue and, 1602

- epicardial and pericoronary fat in patients undergoing coronary angiography, relation to coronary artery calcium and, 380
- Atherosclerosis Risk In Communities (ARIC) Study, coronary heart disease risk prediction based on retinal vascular caliber, 58
- Atorvastatin
- comparison of effectiveness of 10 mg versus 80 mg in reducing major cardiovascular events and repeat revascularization in patients with previous percutaneous coronary intervention (Post Hoc Analysis of the Treating to New Targets [TNT] Study), 1312
 - daily (20 mg), for severe high-density lipoprotein cholesterol deficiency in men, comparison of treatment with fenofibrate (200 mg) versus extended-release niacin (2 g) versus, 1341
 - intensive, plasma prostaglandin E2 levels and metalloproteinase-9 activity in patients with non-ST-elevation acute coronary syndrome, 12
 - meta-analysis of head-to-head randomized controlled trials of benefits and risks of rosuvastatin versus, 1654
 - prognosis and risk factors in patients with asymptomatic aortic stenosis and their modulation by, 743
 - 10 mg versus 80 mg dosage effect on biomarkers of inflammation and oxidative stress with metabolic syndrome, 321
 - up-titration of up to 40 mg of in hypercholesterolemic patients at moderately high risk for coronary heart disease, efficacy and safety of ezetimibe added on to atorvastatin (20 mg) versus, 1489
 - up-titration of up to 80 mg of in hypercholesterolemic patients at moderately high risk for coronary heart disease, efficacy and safety of ezetimibe added on to atorvastatin (40 mg) versus, 1495
- Atrial-atrial (A-A) interval, diagnosis of accessory pathway using the His-His (H-H) interval of the first entrained cycle during ventricular overdrive pacing and, 197
- Atrial fibrillation
- in adults following transcatheter closure of atrial septal defects, 902
 - aldosterone synthase gene polymorphism as determinant in patients with heart failure of, 326
 - anticoagulant and anti-inflammatory responses comparison using enoxaparin versus unfractionated heparin for transesophageal echocardiography-guided cardioversion of, 842
 - benefits and mortality of cardiac resynchronization therapy in patients in sinus rhythm versus patients with (Results of the Spanish Atrial Fibrillation and Resynchronization [SPARE] Study), 444
 - catheter ablation for, real-time three-dimensional echocardiography of left atrial volumes and function comparison with persistence of sinus rhythm versus recurrent atrial fibrillation three months later, 847
 - dynamic changes of left ventricular performance and left atrial volume induced in healthy young adults using the Mueller maneuver and implications for, 1557
 - evaluation of catheter ablation of periatrial ganglionic plexi in patients with, 578
 - feasibility and safety of remote-controlled magnetic navigation for ablation of, 1674
 - first episode of, changing trends of hospital utilization in patients after, 568
 - glomerular filtration rate relation to, 1056
 - heart rate control in patients referred for exercise testing with, 704
 - in hypertensive patients with ischemic stroke, echocardiographic and clinical risk factors for, 1348
 - increased inflammatory cell infiltration in the atrial myocardium of patients with, 861
 - long-term risk with symptomatic gastroesophageal reflux disease and esophagitis of, 1207
 - newly-diagnosed after dual-chamber pacemaker implantation in patients without a previous history of atrial fibrillation, frequency, duration, and predictors of, 450
 - or flutter, sinus rhythm restoration in sleep apnea patients with, 709
 - paroxysmal, anatomic approach for ganglionic plexi ablation in patients with, 330
 - permanent, echo-guided cardiac resynchronization pacing usefulness in patients undergoing "ablate and pace" therapy for, effects of heart rate regularization and left ventricular resynchronization on, 854
 - postoperative, dofetilide as routine prophylaxis for, 506 (Readers' Comments)
 - psoriasis, inflammation, and, 647 (Readers' Comments)
 - transition from paroxysmal to persistent, P-wave dispersion in standard twelve-lead electrocardiography to predict, 573
 - validation of simplified method to determine left atrial volume by computed tomography in patients with, 1567
- Atrial septal defect
- "isolated," in infancy, severe pulmonary hypertension complicating, 340
 - percutaneous, minimally invasive retrieval of a dislodged and impacted device from the abdominal aorta, four months after closure of, 1111
 - transcatheter closure in adults, atrial fibrillation following, 902
- Atropine, early injection in dobutamine stress echocardiography, with versus without chronic beta-blocker therapy in patients with known or suspected coronary heart disease, prognostic value of, 1291
- Balloon dilation, poststent, using a noncompliant balloon versus a semicompliant balloon in intravascular ultrasound evaluation of optimal drug-eluting stent expansion after (from the Poststent Optimal Stent Expansion Trial [POET]), 304
- Bare-metal stents
- diabetes mellitus and, meta-analysis comparison (nine trials) of outcomes with drug-eluting stents versus, 1328
 - drug-eluting stents for narrowing in saphenous vein grafts versus, 530
 - impact of mild or moderate renal insufficiency on intravascular ultrasound analysis of chronic vascular response to paclitaxel-eluting stents and (from the TAXUS IV, V, and VI Trials), 1009
 - indications and complications of drug-eluting stents versus, 32 (Editor's Roundtable)
 - stent thrombosis, myocardial infarction, and mortality after drug-eluting stent coronary intervention in patients with diabetes mellitus versus, 165
 - three months after implantation, neointimal coverage by optical coherence tomography of sirolimus-eluting stents versus, 27
 - two-year clinical outcome with drug-eluting stents in a real-world registry of unprotected left main coronary artery stenosis versus (from the Italian Society of Invasive Cardiology), 1463
- Bariatric surgery, for obesity, cardiovascular risk after, 930
- Beta adrenergic blocking agents (beta blockers)
- cardiac event risk in patients with asthma and long-QT syndrome treated with beta₂ agonists, 866
 - chronic, in patients with known or suspected coronary heart disease, prognostic value of dobutamine stress echocardiography with early injection of atropine with versus without, 1291
 - digoxin and clinical outcomes in systolic heart failure patients on, 1356
 - dosing before, during, and after hospitalization for heart failure (from Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure), 1524
 - failure to prescribe acutely in patients with sustained ventricular tachycardia and ventricular fibrillation following acute myocardial infarction, relation of mortality to (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry), 1427
 - heart rate reduction as mechanism for outcome improvement in patients with systolic chronic heart failure, 506 (Readers' Comments)
 - hypertension treated with, new-onset risk of diabetes mellitus and, 242 (Readers' Comments), *Reply*, 244
 - impact of β 1-adrenergic receptor polymorphisms on susceptibility to heart failure, arrhythmogenesis, prognosis, and response to, 726
 - late mortality in hospitalized patients >70 years of age with systolic heart failure receiving versus not receiving, 1711
 - mortality in patients with implantable cardioverter-defibrillators and, 77
 - survival in patients with severe mitral regurgitation and normal left ventricular ejection fraction and therapy with, 611
- Bifurcation lesions, long-term outcomes of percutaneous coronary intervention of narrowings after, 404
- Biomarkers. *See also* Inflammation, biomarkers
- cardiac, release in long-distance runners aged \geq 55 years, magnitude, reproducibility, and association with baseline cardiac function of, 218
 - of cardiomyocyte damage, frequency of elevation in otherwise healthy newborns of, 761
 - of inflammation and oxidative stress with metabolic syndrome, effect of 10 mg versus 80 mg atorvastatin on, 321

- N-terminal pro-B-type natriuretic peptide, for carcinoid heart disease, 938
- pregnancy associated plasma protein A, in early phase of acute coronary syndrome, 954 (Readers' Comments)
- proteinuria as, among patients undergoing percutaneous coronary intervention of cardiovascular events (data from the Evaluation of Oral Xemilofiban in Controlling Thrombotic Events [EXCITE] Trial), 1151
- in thalassemia intermedia, relation between iron-overload indices, cardiac echo-Doppler and, 363
- Bivalirudin**
- in percutaneous coronary intervention for stable or unstable angina pectoris, bleeding risk and outcomes of glycoprotein IIb/IIIa inhibitors with targeted low-dose unfractionated heparin versus, 160
 - with provisional glycoprotein IIb/IIIa inhibitors in acute myocardial infarction patients complicated by cardiogenic shock undergoing primary angioplasty, 287
- Blacks.** *See also* African Americans
- comparison of cardiovascular risk factors for high brachial pulse pressure in whites versus (Charleston Heart Study, Evans County Study, NHANES I and II Studies), 1514
- Bleeding.** *See also* Hemorrhage
- compliance with prolonged dual antiplatelet therapy (aspirin + thienopyridine) following drug-eluting stent implantation, and incidence of, 1477
 - "nuisance," and clopidogrel compliance in patients undergoing intracoronary drug-eluting stent implantation, 1614
 - in patients undergoing coronary stenting on dual oral antiplatelet treatment, oral anticoagulant therapy and, 1618
 - in patients with recent cerebrovascular or coronary ischemic events, relation between aspirin dose, all-cause mortality, and (from the BRAVO Trial), 1285
 - risk of, and outcomes of bivalirudin versus glycoprotein IIb/IIIa inhibitors with targeted low-dose unfractionated heparin in percutaneous coronary intervention for stable or unstable angina pectoris, 160
- Blood pressure**
- high brachial, comparison of cardiovascular risk factors in blacks versus whites for (Charleston Heart Study, Evans County Study, NHANES I and II Studies), 1514
 - hypertensive, usefulness of response during single-stage exercise testing to predict long-term outcome in patients with peripheral arterial disease of, 921
 - increased paradoxical systolic, after exercise, as predictor of cardiovascular mortality, 518
 - systolic
 - in double product reserve, Duke Treadmill Score nomogram in men adjusted for age and, 1407
 - survival in both ischemic and nonischemic systolic heart failure and, 1698
- Blood transfusion, in acute myocardial infarction, mortality and, 1116 (Readers' Comments)**
- Blood volume, characteristics in anemic patients with low versus preserved left ventricular ejection fractions of, 1069**
- BNP.** *See* Natriuretic peptides, B-type
- Body mass index**
- impact on clinical outcomes in patients with ST-segment elevation acute myocardial infarction, waist-to-hip ratio and (from the Korean Acute Myocardial Infarction Registry), 957
 - mortality in patients with intermediate to high risk of coronary artery disease and, 1028
 - waist circumference in diagnosing metabolic syndrome and identifying apparently healthy subjects at increased risk of cardiovascular disease versus, 40
- Body surface maps**
- for acute myocardial infarction diagnosis, comparison of value of leads from 12-lead electrocardiogram to, 257
 - in patients with acute myocardial infarction, posterior thorax data and, 1117 (Readers' Comments)
- Body weight, limitations of United States cardiac catheterization laboratories for the morbidly obese, 285**
- Bogalusa Heart Study, parental coronary artery disease and adverse effects of metabolic syndrome and aging on carotid artery intima-media thickness, 180**
- BRAVO Trial, relation between aspirin dose, all-cause mortality, and bleeding in patients with recent cerebrovascular or coronary ischemic events, 1285**
- Brugada-type electrocardiograms, prognosis of subjects in a population of middle-aged Japanese diagnosed during a health examination with, 584**
- Bundle branch block, in patients with heart failure, 507 (Readers' Comments)**
- CABG.** *See* Coronary artery bypass grafting
- CAD.** *See* Coronary artery disease
- Calcium, coronary artery**
- and abdominal aortic calcium, indexes of kidney function and (from the Framingham Offspring Study), 440
- C-reactive protein in asymptomatic women with systemic lupus erythematosus or rheumatoid arthritis and, 755**
- defining normal distributions in women and men of (from the Framingham Heart Study), 1136
 - epicardial and pericoronary fat in patients undergoing coronary angiography, relation to coronary atherosclerosis and, 380
- Carcinoid heart disease, usefulness of N-terminal pro-brain natriuretic peptide as biomarker for, 938**
- Cardiac catheterization laboratories**
- quantitative automated assessment of myocardial perfusion in, 980
 - in the United States for, body weight limitations for the morbidly obese in, 285
- Cardiac death, sudden.** *See also* Mortality
- frequency of, after invasive coronary revascularization in acute myocardial infarction, 1
 - prognostic significance of increased left ventricular mass index in patients with stable coronary heart disease to mortality and (from the Heart and Soul Study), 1131
- Cardiac events.** *See also* Cardiovascular events
- future adverse, relation of increased chromosomal damage in patients with known coronary artery disease to, 1296
 - major adverse
 - serum uric acid levels in patients with suspected or definite coronary artery disease and, 797
 - risk in patients with asthma and long-QT syndrome treated with beta₂ agonists, 866
- Cardiac magnetic resonance.** *See* Magnetic resonance imaging, cardiac
- Cardiac resynchronization therapy**
- dual-site left ventricular, 1687
 - echo-guided pacing usefulness in patients undergoing "ablate and pace" therapy for permanent atrial fibrillation, effects of heart rate regularization and left ventricular resynchronization on, 854
 - emerging role of myocardial perfusion imaging to evaluate patients for, 211
 - guided by echocardiography versus guided by electrocardiographic QRS interval width, interventricular delay interval optimization in, 1373
 - long-term, effects of interruption on left ventricular function and dyssynchrony, 718
 - measures of left ventricular dyssynchrony and the correlation to clinical and echocardiographic response after, 598
 - in patients with atrial fibrillation versus patients in sinus rhythm, comparison of benefits and mortality of (Results of the Spanish Atrial Fibrillation and Resynchronization [SPARE] Study), 444
- Cardiac tamponade, Empress Sissi and, 1278**
- Cardiogenic shock, bivalirudin with provisional glycoprotein IIb/IIIa inhibitors in acute myocardial infarction patients undergoing primary angioplasty complicated by, 287**
- Cardiomyocyte apoptosis, right ventricular, acute myocardial infarction of left ventricular wall and, 658**
- Cardiomyocyte damage markers, frequency of elevation in otherwise healthy newborns of, 761**
- Cardiomyopathy**
- arrhythmogenic right ventricular, clinical features in men versus women of, 1252
 - hypertrophic (*See* Hypertrophic cardiomyopathy)
 - idiopathic dilated
 - in children, relation between left ventricular regional radial function and radial wall motion abnormalities using two-dimensional speckle tracking in, 335

- exercise-induced changes of functional mitral regurgitation in asymptomatic or mildly symptomatic patients with, 481
- ischemic
- comparison between tissue Doppler imaging and velocity-encoded magnetic resonance imaging for measurement of myocardial velocities, assessment of left ventricular dyssynchrony, and estimation of left ventricular filling pressures in patients with, 1366
 - comparison of the initial microvolt T-wave alternans screening with repeat testing and long-term prognosis for, 1114 (Readers' Comments)
 - usefulness of microvolt T-wave alternans to predict outcomes beyond one year in patients with, 280
- ischemic or idiopathic dilated
- predicting effects of exercise training in patients with heart failure secondary to, 1073
 - relation of serum heat shock protein 60 level to severity and prognosis in chronic heart failure secondary to, 606
- restrictive, comparison of usefulness of tissue Doppler imaging versus brain natriuretic peptide for differentiation of constrictive pericardial disease from, 357
- Cardiorespiratory fitness, inflammatory markers, fibrinolytic factors, and lipoprotein(a) in patients with type 2 diabetes mellitus and, 700
- Cardiovascular disease
- body mass index versus waist circumference in diagnosing metabolic syndrome identifying apparently healthy subjects at increased risk of, 40
 - caffeinated coffee consumption, heart valve disease in the elderly and (from the Framingham Heart Study), 1502
 - chronic kidney disease as predictor of (from the Framingham Heart Study), 47
 - comparison of three treatment approaches to decreasing risk in nondiabetic insulin-resistant dyslipidemic subjects, 64
 - diabetic, translation between endocrinology and cardiology and, 1115 (Readers' Comments)
 - endothelial dysfunction in, 418 (Editor's Roundtable)
 - endothelial function relation to cardiovascular risk in women with sedentary occupations and without known, 348
 - homozygous familial hypercholesterolemia and, longitudinal evaluation and assessment of, 1438
 - in human immunodeficiency virus-infected patients, 635
 - hypovitaminosis D prevalence in (from the National Health and Nutrition Examination Survey 2001 to 2004), 1540
 - nonsteroidal antiinflammatory drugs and risk of, 1046 (Editor's Roundtable)
 - number of metabolic syndrome risk factors and mortality due to all-causes or, 689
 - and poverty in India, 102
 - reducing residual risk of, lipid management beyond statins and, 559 (Editor's Roundtable)
 - reflections on the utility of imaging for prevention of, 1116 (Readers' Comments)
 - risk
 - stress echocardiography after percutaneous coronary intervention for stratification of, 1170
 - risk, discordance between use of evidence-based medical therapy and physicians' estimation of, 1142
 - risk after bariatric surgery for obesity of, 930
 - self-reported periodontal disease, usefulness to identify individuals with elevated inflammatory markers at risk of, 1509
- Cardiovascular events. *See also* Cardiac events
- major, in patients with previous percutaneous coronary intervention, effectiveness of atorvastatin 10 mg versus 80 mg in reducing (Post Hoc Analysis of the Treating to New Targets [TNT] Study), 1312
 - proteinuria usefulness as prognostic marker among patients undergoing percutaneous coronary intervention of (data from the Evaluation of Oral Xemilofiban in Controlling Thrombotic Events [EXCITE] Trial), 1151
 - usefulness of N-terminal pro-brain natriuretic peptide to predict cardiovascular outcomes in patients with heart failure and preserved left ventricular ejection fractions and, 733
- Cardiovascular imaging
- computed tomographic angiography as model for defining optimal research study design for, 943 (Editorial)
 - for prevention of coronary disease, reflections on the utility of, 1116 (Readers' Comments)
- Cardioverter-defibrillators, implantable
- cost of preimplantation cardiac imaging in patients referred for a primary-prevention, 588
 - in heart failure patients, sleep-disordered breathing impact on life-threatening ventricular arrhythmia with, 1064
 - management of subacute and delayed right ventricular perforation with a lead for, 1352
 - mortality and beta blockers, angiotensin-converting enzyme inhibitors or angiotensin receptor blockers, and statins in patients with, 77
 - role in methadone-induced long-QT syndrome, 370 (Readers' Comments)
- Carnitine supplementation, flow-mediated dilation and vascular inflammatory responses to a high-fat meal in healthy young adults and, 1413
- Carotid artery intima-media thickness
- C-reactive protein in Taiwanese with familial hypercholesterolemia and, 184
 - Framingham risk score for calcium level in coronary arteries or aorta versus, 825
 - parental coronary artery disease and adverse effects of metabolic syndrome and aging on (from the Bogalusa Heart Study), 180
- Catheter ablation, of periatrial ganglionic plexi in patients with atrial fibrillation, 578
- Centers for Medicare and Medicaid Services, chest pain center accreditation and core measures for acute myocardial infarction of, 120
- Charleston Heart Study, comparison of cardiovascular risk factors for high brachial pulse pressure in blacks versus whites, 1514
- CHD. *See* Coronary heart disease
- Chest pain
- non-specific, cardiovascular risk in women with (from the Women's Health Initiative Hormone Trials), 693
 - prognostic significance of troponin T elevation in patients without, 668
 - of suspected cardiac origin, cost-efficiency of myocardial contrast echocardiography in patients presenting with a nondiagnostic electrocardiogram to the emergency department and, 649
- Chest pain centers, Centers for Medicare and Medicaid Services core measures for acute myocardial infarction and accreditation of, 120
- Children. *See also* Congenital heart disease
- comparative outcome of bidirectional Glenn shunt with pulmonary vascular resistance ≥ 3.5 Woods units versus < 3.5 Woods units of, 907
 - with congenital heart disease, managed ventricular pacing in, 866
 - with hypertrophic cardiomyopathy
 - prognosis for, 1418
 - significance of left atrial volume in, 1390
 - with idiopathic dilated cardiomyopathy, relation between left ventricular regional radial function and radial wall motion abnormalities using two-dimensional speckle tracking in, 335
 - long-term risk of fatal malignancy following radiofrequency ablation in, 913
 - long-term safety and efficacy of low-density lipoprotein apheresis for homozygous familial hypercholesterolemia in, 1199
 - transcatheter closure of patent ductus arteriosus using the swivel-disk device versus plug occluder in, 486
 - with ventricular septal defect, exercise capacity and participation of, 1079
- Cholesterol
- high-density lipoprotein (HDL)
 - change in apparently healthy male physicians, incident coronary heart disease and, 1663
 - severe deficiency in men, comparison of treatment with daily atorvastatin (20 mg) versus fenofibrate (200 mg) versus extended-release niacin (2 g) for, 1341
 - low-density lipoprotein
 - apheresis in childhood for homozygous familial hypercholesterolemia, long-term safety and efficacy of, 1199
 - in clinical diagnosis for familial hypercholesterolemia, genetic testing versus, 1187
 - oxidized, oxidative stress and, 1118 (Readers' Comments)
 - treating, in the general population, 372 (Readers' Comments)
- Cholestyramine, in the Post Coronary Artery Bypass Graft (Post CABG) Trial, pleiotropic effect of lovastatin, with and without, 1023

- Chromosomal damage, increased, in patients with known coronary artery disease, future adverse cardiac events and, 1296
- Chronic kidney disease. *See* Kidney disease, chronic
- Chronic obstructive pulmonary disease, statin therapy effect on mortality in patients with peripheral arterial disease with versus without associated, 192
- Chronic total occlusions, "true," comparison of safety, efficacy, and outcome of successful versus unsuccessful percutaneous coronary interventions in, 1175
- Clopidogrel
 - benefit doubt for acute coronary syndrome and percutaneous coronary interventions due to rebound adverse events, 248 (Readers' Comments)
 - comparison of higher loading and maintenance dose to standard dose on platelet function and outcomes after percutaneous coronary intervention using drug-eluting stents, 401
 - compliance in patients undergoing intracoronary drug-eluting stent implantation, impact of "nuisance" bleeding on, 1614
 - loading dose in healthy volunteers, platelet functional recovery dynamics after, 790
 - nonresponsiveness in patients with acute myocardial infarction, reloading effectiveness to overcome, 524
 - plus combined aspirin therapy in diabetes mellitus with coronary heart disease and impaired antiplatelet response to low-dose aspirin, comparison of increased aspirin dose versus, 396
 - short (<1 year) and long-term (≥ 1 year), after percutaneous coronary intervention, comparative impact on mortality of, 1159
 - 600-mg loading dose of, impact on 30-day outcome in unselected patients undergoing percutaneous coronary intervention, 1318
- Cocaine, myocardial infarction and prevalent use of, 966
- Cockcroft-Gault equation, mortality prediction in patients admitted for exclusion of myocardial ischemia with Modification of Diet in Renal Disease versus, 140
- Coffee, caffeinated, consumption of, cardiovascular disease, and heart valve disease in the elderly and (from the Framingham Heart Study), 1502
- Computed tomography. *See also* Coronary angiography
 - angiography, as model for defining optimal research study design for cardiovascular imaging, 943 (Editorial)
 - contrast-enhanced 64-slice, to assess the significance of angiographically ambiguous coronary narrowings, comparison of intravascular ultrasound to, 994
 - multidetector
 - contrast-induced nephropathy incidence in chronic renal insufficiency patients undergoing treated with preventive measures, 353
 - in individuals without known coronary artery disease, costs and clinical outcomes of, from an analysis of Medicare Category III transaction codes, 672
 - multislice
 - clinical significance of epicardial fat measured using, 767
 - coronary atherosclerosis detected by, association of traditional risk factor assessment with, 316
 - 64-slice multidetector
 - for characterizations of noncalcified coronary arterial plaques, spectral analysis of intravascular ultrasound backscatter signals compared with, 988
 - validation of simplified method to determine left atrial volume in patients with atrial fibrillation by, 1567
- Congenital heart disease. *See also* Hypertrophic cardiomyopathy
 - complex, in adults with ventricular dysfunction (subaortic right ventricle and single ventricle physiology) and, prevalence and determinants of, 625
 - erectile dysfunction in men with, 1728
 - hybrid intraoperative pulmonary artery stent placement for, 1737
 - managed ventricular pacing in pediatric patients and patients with, 866
 - physical activity, cardiac function, exercise capacity and quality of life with systemic right ventricle, 1258
- Constrictive pericardial disease, tissue Doppler imaging versus brain natriuretic peptide for differentiation of restrictive cardiomyopathy from, 357
- Continuous quality improvement program, major morbidity after cardiac surgery and, 772
- Contrast agents. *See also* Nephropathy, contrast-induced
 - angiographically ambiguous narrowings of coronary arteries, intravascular ultrasound to assess significance of, compared to 64-slice computed tomography enhanced by, 994
 - in direct percutaneous coronary intervention for ST-elevation myocardial infarction, emergency pretreatment for allergy to, 1469
 - safety of administration in stress echocardiography for endocardial enhancement versus noncontrast stress echocardiography, 1444
 - ultrasound
 - echocardiography with and without, acute mortality in hospitalized patients undergoing (Multicenter Registry Results in 4,300,966 Consecutive Patients), 1742
 - safety in stress echocardiography, 1269
- Controlled Abciximab and Device Investigation to Lower Late Angioplasty complications (CADILLAC), predictive value of, in primary percutaneous coronary intervention for outcomes in ST-elevation acute myocardial infarction comparison with three different risk scores, 6
- Coronary angiography
 - computed tomographic, as model for defining optimal research study design for cardiovascular imaging, 943 (Editorial)
 - diagnostic safety during uninterrupted therapeutic warfarin treatment of, 386
 - epicardial and pericoronary fat relation to coronary atherosclerosis and coronary artery calcium in patients undergoing, 380
 - intravascular ultrasonic detection of myocardial bridging of the left anterior descending coronary artery compared to, 1608
 - multidetector computed tomographic
 - contrast-induced nephropathy incidence in chronic renal insufficiency patients undergoing treated with preventive measures, 353
 - in individuals without known coronary artery disease, costs and clinical outcomes of, from an analysis of Medicare Category III transaction codes, 672
 - myocardial perfusion imaging role in patients with end-stage renal disease undergoing, 1451
 - normal, clinical and electrocardiographic correlates in patients referred for primary percutaneous coronary intervention of, 155
 - routine periodic fasting usefulness to lower risk of coronary artery disease in patients undergoing, 814
 - standard versus three-dimensional, comparison of assessment of native coronary arteries by, 272
 - of vulnerable plaques, 1644 (Editor's Roundtable)
- Coronary angioplasty. *See also* Percutaneous coronary intervention(s)
 - primary, for acute myocardial infarction complicated by cardiogenic shock, bivalirudin with provisional glycoprotein IIb/IIIa inhibitors in patients undergoing, 287
 - primary percutaneous, relation of left ventricular infarct transmural and infarct size after, to time from symptom onset to balloon inflation in, 1163
- Coronary arteries
 - angiographically ambiguous narrowings of, comparison of contrast-enhanced 64-slice computed tomography and intravascular ultrasound to assess significance of, 994
 - assessment of culprit and remote narrowings with long-term outcomes using optical coherence tomography, 391
 - calcium level, Framingham risk score for calcium level in aorta or carotid artery intima-media thickness versus, 825
 - left anterior descending
 - comparison of angiographic versus intravascular ultrasonic detection of myocardial bridging of, 1608
 - hypertrophic cardiomyopathy and prognostic implications of coronary flow reserve in, 1718
 - native, standard versus three-dimensional coronary angiography for assessment of, 272
 - small vessel lesions treated with paclitaxel-eluting stents, clinical and angiographic follow-up of (from the TRUE Registry), 1002
- Coronary artery bypass grafting
 - isolated, impact of early and delayed stroke on in-hospital and long-term mortality after, 411
 - with multivessel coronary artery disease before elective vascular surgery for abdominal aortic and peripheral occlusive disease, 809

- in patients without angina pectoris, prognostic significance of myocardial ischemia by dobutamine stress echocardiography after, 1156
- perioperative myocardial infarction impact on angiographic and clinical outcomes following (from PProject of Ex-vivo Vein graft ENgineering via Transfection [PREVENT] IV), 546
- pleiotropic effect of lovastatin, with and without cholestyramine, in the Post Coronary Artery Bypass Graft (Post CABG) Trial, 1023
- previous, outcomes of patients with acute coronary syndrome and (from the Pravastatin or Atorvastatin Evaluation and Infection Therapy [PROVE IT-TIMI 22] and the Aggrastat to Zocor [A to Z] Trials), 552
- as primary revascularization in patients with acute coronary syndrome, percutaneous coronary intervention versus, 173
- in type 2 diabetes patients, intravenous glucagon-like peptide-1 effects on glucose control and hemodynamics after, 646 (Readers' Comments)
- Coronary artery disease**
- anemia and chronic kidney disease effectiveness as predictor for presence and severity in patients undergoing stress myocardial perfusion study of, 266
- association between serum uric acid levels and perioperative and late cardiovascular outcome in patients undergoing elective vascular surgery with suspected or definite, 797
- costs and clinical outcomes of coronary multidetector computed tomographic angiography in individuals without, from an analysis of Medicare Category III transaction codes, 672
- diabetes mellitus and hypothalamic hypoestrogenemia in premenopausal women and (from the National Heart, Lung, and Blood Institute Sponsored WISE Study), 150
- intermediate to high risk of, relation of exercise capacity and body mass index to mortality in patients with, 1028
- known, relation of increased chromosomal damage to future adverse cardiac events in patients with, 1296
- multivessel, usefulness of revascularization before elective vascular surgery for abdominal aortic and peripheral occlusive disease in patients with, 809
- parental, effect on adverse effects of metabolic syndrome and aging on carotid artery intima-media thickness (from the Bogalusa Heart Study), 180
- peripheral arterial response compared to mental stress in men versus women with, 970
- psoriasis and, 1631 (Editor's Consensus)
- routine periodic fasting usefulness to lower risk in patients undergoing coronary angiography of, 814
- two-year outcome of patients treated with sirolimus- versus paclitaxel-eluting stents in an unselected population with (from the REWARDS Registry), 292
- Coronary flow reserve, prognostic implications on left anterior descending coronary artery in hypertrophic cardiomyopathy of, 1718**
- Coronary heart disease**
- adult-onset asthma relation to stroke and, 508 (Readers' Comments)
- with diabetes mellitus and impaired antiplatelet response to low-dose aspirin, comparison of increased aspirin dose versus combined aspirin plus clopidogrel therapy in, 396
- ezetimibe efficacy and safety added on to atorvastatin (20 mg) versus uptitration of atorvastatin (to 40 mg) in hypercholesterolemic patients at moderately high risk for, 1489
- ezetimibe efficacy and safety added on to atorvastatin (40 mg) versus uptitration of atorvastatin (to 80 mg) in hypercholesterolemic patients at moderately high risk for, 1495
- incident, in apparently healthy male physicians, change in high-density lipoprotein cholesterol and, 1663
- increased prevalence in US adults without chronic kidney disease, serum cystatin C and, 54
- left atrial volume index usefulness to predict heart failure hospitalization and mortality in ambulatory patients with, comparison to left ventricular ejection fraction and (from the Heart and Soul Study), 70
- prognostic value of dobutamine stress echocardiography with early injection of atropine with versus without chronic beta-blocker therapy in patients with known or suspected, 1291
- risk prediction, retinal vascular caliber and (from the Atherosclerosis Risk In Communities [ARIC] Study), 58
- self-reported leisure-time physical activity usefulness to predict long-term survival in patients with, 375
- stable, prognostic significance of increased left ventricular mass index to mortality and sudden death in patients with (from the Heart and Soul Study), 1131
- usefulness of fasting plasma glucose to predict mortality in persons ≥ 60 years of age without diabetes mellitus or in those with undiagnosed diabetes mellitus (from The Dubbo Study), 825
- Coronary sinus, mitral valve annulus and, influence of left atrial and ventricular volumes on, 890**
- C-reactive protein**
- carotid artery intima-media thickness in Taiwanese with familial hypercholesterolemia and, 184
- coronary artery calcium in asymptomatic women with systemic lupus erythematosus or rheumatoid arthritis and, 755
- differential impact of admission levels in patients with ST-elevation versus non-ST-elevation myocardial infarction on 28-day mortality risk of (from the monitoring trends and determinants on cardiovascular diseases [MONICA]/Cooperative Health Research in the Region of Augsburg [KORA] Augsburg Myocardial Infarction Registry), 1125
- epidemiology, heritability, and genetic linkage in African Americans of (from the Jackson Heart Study), 835
- serum levels in patients with microalbuminuria, foscipril treatment and, 223
- Creatine kinase, MB level (infarct size) in patients presenting with acute myocardial infarction, relation between previous lipid-lowering therapy, 1119**
- Critically ill patients, without acute coronary syndrome, prognostic implications of normal (<0.10 ng/ml) and borderline (0.10 to 1.49 ng/ml) troponin elevation levels in, 509**
- CRT. See Cardiac resynchronization therapy**
- CRUSADE National Quality Improvement Initiative, patterns of upstream antiplatelet therapy before primary percutaneous coronary intervention for acute ST-elevation myocardial infarction, 1335**
- Cystatin C**
- clinical correlates and heritability of (from the Framingham Offspring Study), 1194
- serum levels and increased prevalence of coronary heart disease in US adults without chronic kidney disease, 54
- Dallas Heart Study, association between renal function and circulating levels of natriuretic peptides, 1394**
- Death. See Cardiac death, sudden; Mortality**
- Defibrillators. See Cardioverter-defibrillators**
- Developing countries, native valve infective endocarditis in a tertiary care center in Tunisia, 1247**
- Diabetes mellitus**
- cardiovascular disease and, translation between endocrinology and cardiology and, 1115 (Readers' Comments)
- cardiovascular mortality in patients older than 65 years and, 1668
- with coronary heart disease and impaired antiplatelet response to low-dose aspirin, comparison of increased aspirin dose versus combined aspirin plus clopidogrel therapy in, 396
- fasting plasma glucose usefulness to predict mortality of coronary heart disease in persons ≥ 60 years of age with undiagnosed or without (from The Dubbo Study), 825
- hypertension treated with beta blockers, new-onset risk of, 242 (Readers' Comments), *Reply*, 244
- hypothalamic hypoestrogenemia and coronary artery disease in premenopausal women and (from the National Heart, Lung, and Blood Institute Sponsored WISE Study), 150
- intravascular ultrasound assessment of expansion of the sirolimus-eluting (Cypher Select) and paclitaxel-eluting (Taxus Express-2) stent in patients with, 19
- meta-analysis comparison (nine trials) of outcomes with drug-eluting stents versus bare metal stents in patients with, 1328
- normal myocardial perfusion imaging and heart rate response to adenosine in patients with, 1103
- stent thrombosis, myocardial infarction, and mortality after drug-eluting versus bare-metal stent coronary intervention in patients with, 165

- type 2
 cardiorespiratory fitness relation to inflammatory markers, fibrinolytic factors, and lipoprotein(a) in patients with, 700
 efficacy and safety of a potent and selective peroxisome proliferator activated receptor alpha agonist in subjects with dyslipidemia and, 434
 endothelial dysfunction in impaired fasting glycemia, impaired glucose tolerance, and, 497
 intravenous glucagon-like peptide-1 effects on glucose control and hemodynamics after coronary artery bypass surgery in patients with, 646 (Readers' Comments)
- Digoxin
 clinical outcomes in systolic heart failure patients on contemporary background heart failure therapy and, 1356
 reduction of heart failure hospitalization in chronic systolic and diastolic heart failure and, 1681
- Dofetilide, as routine prophylaxis for postoperative atrial fibrillation prevention after coronary surgery, 506 (Readers' Comments)
- Doppler imaging. *See* Tissue Doppler imaging
- Drug-eluting stents. *See* Stents, drug-eluting
- The Dubbo Study, usefulness of fasting plasma glucose to predict mortality of coronary heart disease in persons ≥ 60 years of age without diabetes mellitus or in those with undiagnosed diabetes mellitus, 825
- Duke Treadmill Score, age and double product (systolic blood pressure \times heart rate) reserve-adjusted modification of the nomogram in men, 1407
- Dyslipidemia
 efficacy and safety of a potent and selective peroxisome proliferator activated receptor alpha agonist in subjects with type 2 diabetes mellitus and, 434
 mixed, combined statin plus omega-3 fatty acid therapy for, 1040
 mixed, effects of adding prescription omega-3 acid ethyl esters to simvastatin (20 mg/day) on lipids and lipoprotein particles in men and women with, 429, 1425 (Correction)
 nondiabetic insulin-resistant, comparison of three treatment approaches to decreasing cardiovascular disease risk in, 64
- Dyspnea, as reason for exercise testing termination in patients with heart failure, prognostic usefulness of fatigue versus, 879
- Ebstein anomaly, operation for, left ventricular function and, 1724
- Echocardiography. *See also* Tissue Doppler imaging; Ultrasound
 after cardiac resynchronization therapy, correlation of left ventricular dyssynchrony measures to clinical response and, 598
 dobutamine stress
 after coronary revascularization, prognostic significance of myocardial ischemia in patients without angina pectoris and, 1156
 with early injection of atropine with versus without chronic beta-blocker therapy in patients with known or suspected coronary heart disease, prognostic value of, 1291
 in men with acutely decompensated chronic heart failure, clinical and neurohumoral improvement prediction after levosimendan treatment and, 1225
- Doppler, in patients with bileaflet mechanical prostheses or bioprosthesis in the aortic valve position, discrepancies with catheter gradients and, 1383
- interventricular delay interval optimization in cardiac resynchronization therapy guided by electrocardiographic QRS interval width versus, 1373
- myocardial contrast, in patients presenting to the emergency department with chest pain of suspected cardiac origin and a nondiagnostic electrocardiogram, cost-efficiency of, 649
- preimplantation, in patients referred for a primary-prevention implantable cardioverter-defibrillator, cost of, 588
- real-time three-dimensional
 accuracy and reproducibility of left ventricular function quantitation by cardiac magnetic resonance versus, 778
 left atrial volumes and function in patients having catheter ablation for atrial fibrillation comparison with persistence of sinus rhythm versus recurrent atrial fibrillation three months later, 847
 to predict left ventricular remodeling after acute myocardial infarction, left ventricular conic index measured by, 1433
 quantitative analysis of left atrial function in left ventricular diastolic dysfunction, 1097
- stress
 noncontrast during, safety of contrast administration for endocardial enhancement compared with, 1444
 for risk stratification of patients after percutaneous coronary intervention, 1170
 safety of ultrasound contrast agents in, 1269
- three-dimensional
 differentiation of hypertrophic cardiomyopathy from other forms of left ventricular hypertrophy with, 616
 new generation of, for right ventricular volumetric and functional measurements, 499
- transesophageal, cardioversion guided by, anticoagulant and anti-inflammatory responses of enoxaparin versus unfractionated heparin for atrial fibrillation in, 842
 with and without an ultrasound contrast agent, acute mortality in hospitalized patients undergoing (Multicenter Registry Results in 4,300,966 Consecutive Patients), 1742
- Editor's Roundtable discussions, informativeness and usefulness of, 507 (Readers' Comments)
- Educational level, inflammation-sensitive biomarker level and, 1034
- Effect of Combination Ezetimibe and High-Dose Simvastatin versus Simvastatin Alone on the Atherosclerotic Process in Patients With Heterozygous Familial Hypercholesterolemia (ENHANCE) trial, public hype and careful analysis of, 368 (Editorial)
- Ejection fractions
 assessment of, cost of preimplantation cardiac imaging in patients referred for a primary-prevention implantable cardioverter-defibrillator and, 588
- left ventricular
 comparison of $>50\%$ in hospitalized heart failure patients versus those with ejection fraction $<50\%$, one-year outcome (death and rehospitalization) in, 79
 left atrial volume index usefulness to predict heart failure hospitalization and mortality in ambulatory patients with coronary heart disease comparison to (from the Heart and Soul Study), 70
 low versus preserved, comparison of blood volume characteristics in anemic patients with, 1069
 normal, effect of beta-blocker therapy on survival in patients with severe mitral regurgitation and, 611
 in patients with heart failure, left ventricular systolic dyssynchrony relation to QRS duration and, 602
 preserved, usefulness of N-terminal pro-brain natriuretic peptide and brain natriuretic peptide to predict cardiovascular outcomes in patients with heart failure and, 733
- Elderly. *See also* Aging
 caffeinated coffee consumption, cardiovascular disease, and heart valve disease in (from the Framingham Heart Study), 1502
- Electrocardiogram(s)
 alcohol septal ablation for hypertrophic cardiomyopathy and patterns of, 621
 Brugada-type, prognosis of subjects in a population of middle-aged Japanese diagnosed during a health examination with, 584
 nondiagnostic, cost-efficiency of myocardial contrast echocardiography in patients presenting with chest pain of suspected cardiac origin to the emergency department and, 649
 in patients with acute myocardial infarction, posterior thorax data and, 1117 (Readers' Comments)
 respiratory artifact in, significance of, 1090
 12-lead, for acute myocardial infarction diagnosis, comparison of value of leads from body surface maps to, 257
- Electrocardiographic QRS interval width, interventricular delay interval optimization in cardiac resynchronization therapy guided by echocardiography versus, 1373
- Electrocardiography
 during exercise testing, unconventional signs of ischemia during, 949
 in patients with suspected reflex neurally-mediated syncope, reproducibility of findings, 1518
 twelve-lead, standard, P-wave dispersion to predict transition from paroxysmal to persistent atrial fibrillation, 573
- Elizabeth, Empress of Austria ("Sissi"), cardiac tamponade and, 1278

- Emergency department
- cost-efficiency of myocardial contrast echocardiography in patients presenting with chest pain of suspected cardiac origin and a nondiagnostic electrocardiogram to, 649
 - disparities in same-day percutaneous coronary intervention for patients with ST-elevation myocardial infarction in Florida, 2001–2005 in, 802
 - illicit stimulant use in a United States heart failure population presenting to (from the Acute Decompensated Heart Failure National Registry Emergency Module), 1216
- Endocarditis, infective, native valve, in a tertiary care center in a developing country (Tunisia), 1247
- Endothelial cell dysfunction
- in cardiovascular disease, 418 (Editor's Roundtable)
 - in impaired fasting glycemia, impaired glucose tolerance, and type 2 diabetes mellitus, 497
- Endothelial cell function
- cardiovascular risk in women with sedentary occupations and without known cardiovascular disease and relation to, 348
 - predictors in employees with sedentary occupations in a worksite exercise program, 820
- Enoxaparin, anticoagulant and anti-inflammatory responses of unfractionated heparin for transesophageal echocardiography-guided cardioversion of atrial fibrillation versus, 842
- Epicardial adipose tissue, coronary atherosclerosis and, 1602
- Erectile dysfunction, in men with congenital heart disease, 1728
- Esophagitis, long-term risk of atrial fibrillation with symptomatic gastroesophageal reflux disease and, 1207
- EuroSCORE, logistic, compared to brain natriuretic peptide plasma levels in predicting in-hospital and late postoperative mortality in patients undergoing aortic valve replacement for symptomatic aortic stenosis, 749
- Evaluation of Oral Ximilofiban in Controlling Thrombotic Events [EXCITE] Trial, proteinuria usefulness as prognostic marker of mortality and cardiovascular events among patients undergoing percutaneous coronary intervention, 1151
- Evans County Study, comparison of cardiovascular risk factors for high brachial pulse pressure in blacks versus whites, 1514
- Everolimus-eluting stents, Xience V, overview of the 2007 Food and Drug Administration Circulatory System Devices Panel Meeting on, 1624
- Evidence-based medicine, discordance between physicians' estimation of cardiovascular risk and use of, 1142
- Exercise
- functional mitral regurgitation changes in asymptomatic or mildly symptomatic patients with idiopathic dilated cardiomyopathy induced by, 481
 - group-based high-intensity aerobic interval-training program in patients with chronic heart failure, long-term effects of, 1220, 1361
 - isometric, left ventricular diastolic function in trained male weightlifters at rest and during, 97
 - paradoxical systolic blood pressure increase, as predictor of cardiovascular mortality after, 518
 - training in patients with heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, predicting effects of, 1073
 - worksite program, predictors of endothelial function in employees with sedentary occupations in, 820
- Exercise capacity
- mortality in patients with intermediate to high risk of coronary artery disease and, 1028
 - and participation of children with a ventricular septal defect, 1079
 - in patients with systemic right ventricle, physical activity, cardiac function, quality of life and, 1258
- Exercise testing
- cardiopulmonary variables in diastolic versus systolic heart failure, 203
 - differences in treadmill tolerance parameters between patients with partial and advanced interatrial depolarization abnormality, 866
 - in heart failure patients, reproducibility of peak oxygen uptake and other cardiopulmonary parameters (from the Heart Failure and A Controlled Trial Investigating Outcomes of exercise training), 712
 - heart rate control in patients with atrial fibrillation referred for, 704
 - single-stage to predict long-term outcome in patients with peripheral arterial disease, usefulness of hypertensive blood pressure response during, 921
 - stress
 - low-doses of alcohol and warm-up phenomenon in patients with stable angina pectoris during, 146
 - in male firefighters, comparison of cardiocirculatory and thermal strain during fire suppression to, 1551
 - termination in patients with heart failure, prognostic usefulness of dyspnea versus fatigue as reason for, 879
 - unconventional electrocardiographic signs of ischemia during, 949
- Ezetimibe
- added on to atorvastatin (20 mg) versus uptitration of atorvastatin (to 40 mg) in hypercholesterolemic patients at moderately high risk for coronary heart disease, efficacy and safety of, 1489
 - added on to atorvastatin (40 mg) versus uptitration of atorvastatin (to 80 mg) in hypercholesterolemic patients at moderately high risk for coronary heart disease, efficacy and safety of, 1495
 - atherosclerotic process in patients with heterozygous familial hypercholesterolemia and (from the ENHANCE trial), public hype and careful analysis of, 368 (Editorial)
 - thrice weekly, effectiveness of, 1205
- Fasting, routine periodic, usefulness to lower risk of coronary artery disease in patients undergoing coronary angiography of, 814
- Fat
- epicardial, measured using cardiac multislice computed tomography, clinical significance of, 767
 - epicardial and pericoronary, in patients undergoing coronary angiography, relation to coronary atherosclerosis and coronary artery calcium of, 380
 - high dietary, in healthy young adults, carnitine supplementation effect on flow-mediated dilation and vascular inflammatory responses to, 1413
- Fatigue, as reason for exercise testing termination in patients with heart failure, prognostic usefulness of dyspnea versus, 879
- Fenofibrate, for severe high-density lipoprotein cholesterol deficiency in men, comparison of treatment with daily atorvastatin (20 mg) versus extended-release niacin (2 g) versus 200 mg of, 1341
- Fibrinolytic factors, cardiorespiratory fitness in patients with type 2 diabetes mellitus and serum levels of, 700
- Fibroatheromas, thin-capped, in patients with acute coronary syndrome, distribution and frequency in the entire culprit coronary artery as determined by optical coherence tomography of, 975
- Firefighters, male, comparison of cardiocirculatory and thermal strain during fire suppression to exercise stress tests and aerobic exercise testing of, 1551
- Food and Drug Administration Circulatory System Devices Panel Meeting (2007) on the Xience V everolimus-eluting stent, overview of, 1624
- Fosinopril, serum C-reactive protein levels in patients with microalbuminuria and, 223
- Framingham Heart Study
- caffeinated coffee consumption, cardiovascular disease, and heart valve disease in the elderly, 1502
 - chronic kidney disease as predictor of cardiovascular disease, 47
 - defining normal distributions of coronary artery calcium in women and men, 1136
- Framingham Offspring Study
- clinical correlates and heritability of cystatin C, 1194
 - indexes of kidney function coronary artery and abdominal aortic calcium, 440
- Framingham risk score
- coronary atherosclerosis detected by multislice tomography and, 316
 - subclinical atherosclerosis evaluated across three arterial sites and, 825
- Ganglionic plexi
- ablation in patients with paroxysmal atrial fibrillation, anatomic approach for, 330
 - periatrial, evaluation of catheter ablation of, in patients with atrial fibrillation, 578
- Gastroesophageal reflux disease, symptomatic, long-term risk of atrial fibrillation with esophagitis and, 1207
- Gender differences. *See also* Men; Women
- in advanced decompensated heart failure, 454

- Genetic testing, clinical diagnosis for familial hypercholesterolemia versus, 1187
- Geographic differences, clinical presentation, treatment, and outcomes in type A acute aortic dissection and (from the International Registry of Acute Aortic Dissection), 1562
- Giant cell myocarditis, usefulness of immunosuppression for, 1535
- Glenn shunt, bidirectional, comparative outcome in patients with pulmonary vascular resistance ≥ 3.5 Woods units versus < 3.5 Woods units of, 907
- Global Registry for Acute Coronary Events (GRACE)
 magnitude and prognosis of ventricular arrhythmias associated in patients hospitalized with acute coronary syndromes, 1577
 predictive value of, in primary percutaneous coronary intervention for outcomes in ST-elevation acute myocardial infarction comparison with three different risk scores, 6
- Glomerular filtration rate, atrial fibrillation relation to, 1056
- Glucagon-like peptide-1, intravenous, effects on glucose control and hemodynamics in type 2 diabetes patients after coronary artery bypass surgery, 646 (Readers' Comments)
- Glucose
 control and hemodynamics, in type 2 diabetes patients after coronary artery bypass surgery, intravenous glucagon-like peptide-1 effects on, 646 (Readers' Comments)
 endothelial dysfunction in impaired fasting glycemia, type 2 diabetes mellitus and impaired tolerance to, 497
 fasting plasma
 ≤ 100 , 101 to 125, and > 125 mg/dl, and myocardial reperfusion in patients undergoing percutaneous coronary intervention for ST-elevation myocardial infarction, 1457
 usefulness to predict mortality or coronary heart disease in persons ≥ 60 years of age without diabetes mellitus or in those with undiagnosed diabetes mellitus (from The Dubbo Study), 825
- Glycemia, impaired fasting, endothelial dysfunction in impaired glucose tolerance, type 2 diabetes mellitus and, 497
- Glycoprotein IIb/IIIa inhibitors
 provisional, bivalirudin in patients with acute myocardial infarction complicated by cardiogenic shock undergoing primary angioplasty and, 287
 retroperitoneal hemorrhage following percutaneous coronary intervention and, 1473
 with targeted low-dose unfractionated heparin in percutaneous coronary intervention for stable or unstable angina pectoris, bleeding risk and outcomes of bivalirudin versus, 160
- HDL cholesterol. *See* Cholesterol, high-density lipoprotein
- Heart and Soul Study
 left atrial volume index versus left ventricular ejection fraction usefulness to predict heart failure hospitalization and mortality in ambulatory patients with coronary heart disease, 70
 prognostic significance of increased left ventricular mass index to mortality and sudden death in patients with stable coronary heart disease, 1131
- Heart disease. *See* Carcinoid heart disease; Congenital heart disease; Coronary heart disease
- Heart failure
 advanced decompensated, gender differences in patients admitted with, 454
 after acute myocardial infarction, statin therapy and risk of, 1706
 alcohol consumption in hypertensive US male physicians and, 593
 aldosterone synthase gene polymorphism as a determinant of atrial fibrillation in patients with, 326
 $\beta 1$ -adrenergic receptor polymorphisms' impact on arrhythmogenesis, prognosis, response to beta-blocker therapy, and susceptibility to, 726
 beta-blocker therapy dosing before, during, and after hospitalization for (from Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure), 1524
 bundle branch block in patients with, 507 (Readers' Comments)
- chronic
 acutely decompensated, dobutamine-induced changes of the two-dimensional longitudinal deformation usefulness to predict clinical and neurohumoral improvement in men after levosimendan treatment in, 1225
 frequency of, after invasive coronary revascularization in acute myocardial infarction, 1
 group-based high-intensity aerobic interval training in patients with, 1361
 long-term effects of a group-based high-intensity aerobic interval-training program in patients with, 1220
 patterns of structural and functional remodeling of the left ventricle in, 459
 secondary to ischemic or idiopathic dilated cardiomyopathy, relation of serum heat shock protein 60 level to severity and prognosis in, 606
 systolic and diastolic, digoxin and reduction of heart failure hospitalization in, 1681
 telomere length and renal dysfunction in, 207
- chronic systolic
 heart rate reduction as mechanism for beta blockade outcome improvement in patients with, 506 (Readers' Comments)
 stabilized by therapy, prognostic value of N-terminal pro-type-B natriuretic peptide and Doppler left ventricular diastolic variables in patients with, 463
 diastolic versus systolic, cardiopulmonary exercise variables in, 203
 dynamic changes of left ventricular performance and left atrial volume induced in healthy young adults using the Mueller maneuver and implications for, 1557
 hospitalization for, exposure to fine particulate air pollution and, 1230
 hospitalized patients with left ventricular ejection fraction $> 50\%$ versus those with ejection fraction $< 50\%$, comparison of one-year outcome (death and rehospitalization) in, 79
 hospital length of stay influence on quality of care for, 1693
 hospital readmission in patients with, and lack of knowledge about dietary sodium to, 1212
 illicit stimulant use in a United States population presenting to the emergency department with (from the Acute Decompensated Heart Failure National Registry Emergency Module), 1216
 with implantable cardioverter-defibrillator, sleep-disordered breathing impact on life-threatening ventricular arrhythmia in, 1064
 left ventricular systolic dyssynchrony relation to left ventricular ejection fraction and to QRS duration in patients with, 602
 N-terminal pro-brain natriuretic peptide prognostic usefulness in patients with versus without chronic kidney disease and with, 469
 predicting effects of exercise training in patients with ischemic or idiopathic dilated cardiomyopathy secondary to, 1073
 prognostic usefulness of dyspnea versus fatigue as reason for exercise test termination in patients with, 879
 reproducibility of peak oxygen uptake and other cardiopulmonary exercise testing parameters in patients with (from the Heart Failure and A Controlled Trial Investigating Outcomes of exercise training), 712
- systolic
 digoxin and clinical outcomes in patients on contemporary background heart failure therapy, 1356
 late mortality in hospitalized patients > 70 years of age receiving versus not receiving beta blockers, 1711
 prognostic value of pulse-wave tissue Doppler parameters in patients with, 722
 relation of systolic blood pressure to survival in both ischemic and nonischemic, 1698
- Heart Failure and A Controlled Trial Investigating Outcomes of exercise training, reproducibility of peak oxygen uptake and other cardiopulmonary exercise testing parameters in patients with heart failure, 712
- Heart rate
 autonomic control at rest and during unloading of the right ventricle in adolescents with repaired tetralogy of Fallot, 1085
 control in atrial fibrillation patients referred for exercise testing, 704
 in double product reserve, Duke Treadmill Score nomogram in men adjusted for age and, 1407
 regularization, echo-guided cardiac resynchronization pacing usefulness in patients undergoing "ablate and pace" therapy for permanent atrial fibrillation, effects of left ventricular resynchronization and, 854
 response to adenosine in patients with diabetes mellitus and normal myocardial perfusion imaging, 1103

- Heart valve disease, caffeinated coffee consumption, cardiovascular disease in the elderly and (from the Framingham Heart Study), 1502
- Heat shock protein 60, serum level, relation to severity and prognosis in chronic heart failure secondary to ischemic or idiopathic dilated cardiomyopathy, 606
- Hemorrhage. *See also* Bleeding
 intracerebral, and subarachnoid, relation of troponin I levels to in-hospital mortality in patients with ischemic stroke and, 632
 retroperitoneal, following percutaneous coronary intervention, risk predictors of, 1473
 subarachnoid, relation of cardiac troponin I to clinical severity, cardiac dysfunction, and pulmonary congestion in patients with, 1545
- Heparin
 targeted low-dose unfractionated, with glycoprotein IIb/IIIa inhibitors in percutaneous coronary intervention for stable or unstable angina pectoris, bleeding risk and outcomes of bivalirudin versus, 160
 unfractionated, anticoagulant and anti-inflammatory responses of enoxaparin for transesophageal echocardiography-guided cardioversion of atrial fibrillation versus, 842
 unfractionated heparin or low-molecular-weight heparin
 perioperative bridging therapy with patients with mechanical prosthetic heart valves on long-term oral anticoagulants with (from the REGIMEN Registry), 879
- Hepatocellular carcinoma, with intracavitary cardiac involvement: a case report and review of the literature, 643
- Highly active antiretroviral therapy, increased cardiovascular risk in patients with human immunodeficiency virus infection under, 373 (Readers' Comments)
- His-His (H-H) interval, diagnosis of accessory pathway using the atrial-atrial (A-A) interval of the first entrained cycle during ventricular overdrive pacing and, 197
- Homocysteine, free plasma levels of, prognostic value in hospitalized acute coronary syndrome patients, 135
- Hospitals
 discharge from, weekend/holiday versus weekday, guideline adherence and (from the American Heart Association's Get With the Guidelines – Coronary Artery Disease Database), 663
 native valve infective endocarditis in a tertiary care center in Tunisia, 1247
 quality of care for heart failure and length of stay in, 1693
 readmission to, in patients with heart failure, lack of knowledge about dietary sodium and, 1212
 safety and efficacy of offsite percutaneous coronary intervention in 1,348 consecutive patients in rural Tasmania, 1323
 time to diagnosis for acute aortic dissection in, 1399
 utilization in patients after first episode of atrial fibrillation, changing trends of, 568
- Human immunodeficiency virus
 cardiovascular manifestations in, 635
 highly active antiretroviral therapy for, increased cardiovascular risk in, 373 (Readers' Comments)
- Hypercholesterolemia
 ezetimibe efficacy and safety added on to atorvastatin (20 mg) versus uptitration of atorvastatin (to 40 mg) in patients at moderately high risk for coronary heart disease with, 1489
 ezetimibe efficacy and safety added on to atorvastatin (40 mg) versus uptitration of atorvastatin (to 80 mg) in patients at moderately high risk for coronary heart disease with, 1495
- familial
 comparison of genetic versus clinical diagnosis in, 1187
 C-reactive protein and carotid artery intima-media thickness in Taiwanese with, 184
 Effect of Combination Ezetimibe and High-Dose Simvastatin versus Simvastatin Alone on the Atherosclerotic Process in Patients With Heterozygous Familial Hypercholesterolemia (ENHANCE) trial, public hype and careful analysis of, 368 (Editorial)
- homozygous familial
 longitudinal evaluation and assessment of cardiovascular disease in patients with, 1438
 long-term safety and efficacy of low-density lipoprotein apheresis in childhood for, 1199
- Hypertension. *See also* Prehypertension
 alcohol consumption and heart failure in US male physicians with, 593
 heart disease with, association of myocardial strain and left ventricular geometry with progression in, 87
 in patients with ischemic stroke, echocardiographic and clinical risk factors for atrial fibrillation and, 1348
 pulmonary, comparative outcome in patients of bidirectional Glenn shunt with resistance of ≥ 3.5 Woods units versus < 3.5 Woods units in, 907
 pulmonary arterial
 evidence for early right ventricular and septal mechanical activation (interventricular dyssynchrony) in, 1273
 severe, complicating "isolated" atrial septal defect in infancy, 340
 treated with beta blockers, new-onset risk of diabetes mellitus and, 242 (Readers' Comments), *Reply*, 244
- Hypertrophic cardiomyopathy
 alcohol septal ablation effect on the electrocardiographic pattern in patients with, 621
 differentiation from other forms of left ventricular hypertrophy with three-dimensional echocardiography, 616
 first patient clinically diagnosed with, 1418
 left atrial volume significance in patients < 20 years of age with, 1390
 prognostic implications of coronary flow reserve on left anterior descending coronary artery in, 1718
- Hypothalamic hypoestrogenemia, diabetes mellitus and coronary artery disease in premenopausal women and (from the National Heart, Lung, and Blood Institute Sponsored WISE Study), 150
- Hypovitaminosis D, prevalence in cardiovascular diseases (from the National Health and Nutrition Examination Survey 2001 to 2004), 1540
- Immunosuppression, for giant cell myocarditis, usefulness of, 1535
- India, poverty and cardiovascular disease in, 102
- Infants
 newborn, elevation of cardiomyocyte damage markers, frequency in otherwise healthy, 761
 severe pulmonary hypertension complicating "isolated" atrial septal defect in, 340
- Inferior vena cava filters, usage by various surgical and medical subspecialists in post-surgical patients, 226
- Inflammation. *See also* Biomarkers
 in the atrial myocardium of patients with atrial fibrillation, 861
 biomarkers
 adhesion molecules in hypertensive and normotensive postmenopausal women, effect of soy nuts on, 84
 cardiorespiratory fitness in patients with type 2 diabetes mellitus and, 700
 circulating levels in patients with peripheral arterial disease, physical activity during daily life and, 1263
 educational level and, 1034
 elevated, self-reported periodontal disease usefulness to identify individuals at risk of cardiovascular disease with, 1509
 in hypertensive and normotensive postmenopausal women, effect of soy nuts on adhesion molecules and, 84
 psoriasis, atrial fibrillation, and, 647 (Readers' Comments)
- Insulin resistance, in nondiabetic dyslipidemic subjects, comparison of three treatment approaches to decreasing cardiovascular disease risk in, 64
- Interatrial block, partial versus advanced, differences in treadmill exercise tolerance parameters between patients with, 866
- International Registry of Acute Aortic Dissection, geographic differences, clinical presentation, treatment, and outcomes in type A acute aortic dissection, 1562
- Interventricular dyssynchrony, in pulmonary arterial hypertension, evidence for early right ventricular and septal mechanical activation, 1273
- Intracardiac left-to-right shunt(s), migraine headache relief after patent foramen ovale closure and, 916
- Iron-overload indices, in thalassemia intermedia, relation between cardiac echodoppler, biochemical markers and, 363

- Ischemia. *See also* Cardiomyopathy, ischemic; Myocardial ischemia; Stroke, ischemic
 unconventional electrocardiographic signs during exercise testing of, 949
 Italian Society of Invasive Cardiology, two-year clinical outcome with drug-eluting stents versus bare-metal stents in a real-world registry of unprotected left main coronary artery stenosis, 1463
- Jackson Heart Study, epidemiology, heritability, and genetic linkage of C-reactive protein in African Americans, 835
- Kidney disease, chronic
 cardiovascular mortality in patients older than 65 years and, 1668
 N-terminal pro–brain natriuretic peptide prognostic usefulness in patients with heart failure and with versus without, 469
 patients on dialysis with acute coronary syndrome and, impact of optimal medical therapy and revascularization on outcome of, 535
 as predictor for presence and severity of coronary artery disease in patients undergoing stress myocardial perfusion study, effectiveness of, 266
 as predictor of cardiovascular disease (from the Framingham Heart Study), 47
 serum cystatin C and increased coronary heart disease prevalence in US adults without, 54
- Kidney function. *See also* Nephropathy
 indexes of coronary artery and abdominal aortic calcium and (from the Framingham Offspring Study), 440
- Korean Acute Myocardial Infarction Registry, body mass index and waist-to-hip ratio impact on clinical outcomes in patients with ST-segment elevation myocardial infarction, 957
- LDL cholesterol. *See* Cholesterol, low-density lipoprotein
- Left atrial function
 in left ventricular diastolic dysfunction, quantitative analysis using real-time three-dimensional echocardiography of, 1097
 real-time three-dimensional echocardiography comparison in patients having catheter ablation for atrial fibrillation with persistence of sinus rhythm versus recurrent atrial fibrillation three months later of, 847
- Left atrial size, left ventricular mass in National Football League linemen versus nonlinemen and, 343
- Left atrial volume
 index of, left ventricular ejection fraction usefulness to predict heart failure hospitalization and mortality in ambulatory patients with coronary heart disease comparison to (from the Heart and Soul Study), 70
 in patients < 20 years of age with hypertrophic cardiomyopathy, significance of, 1390
 in patients with atrial fibrillation, validation of simplified method using computed tomography to determine, 1567
 real-time three-dimensional echocardiography comparison in patients having catheter ablation for atrial fibrillation with persistence of sinus rhythm versus recurrent atrial fibrillation three months later of, 847
 relation between mitral valve annulus and coronary sinus, influence of left ventricular volume and, 890
- Left ventricle
 dual-site cardiac resynchronization therapy on, 1687
 structural and functional remodeling patterns in chronic heart failure, 459
 usefulness of shape of, to predict the early recovery of left ventricular function after isolated aortic valve replacement for aortic valve stenosis, 1530
- Left ventricular conic index, measured by real-time three-dimensional echocardiography to predict left ventricular remodeling after acute myocardial infarction, 1433
- Left ventricular dysfunction. *See also* Left ventricular function
 diastolic, quantitative analysis using real-time three-dimensional echocardiography of left atrial function in, 1097
 drug-eluting stent safety in patients undergoing percutaneous coronary intervention with, 679
- Left ventricular dyssynchrony
 comparison by two-dimensional speckle tracking versus tissue Doppler imaging in patients with non–ST-elevation myocardial infarction and preserved left ventricular systolic function, 1146
 effects of interruption of long-term cardiac resynchronization therapy on left ventricular function and, 718
 measures of, and correlation to clinical and echocardiographic response after cardiac resynchronization therapy, 598
 in patients with non–ST-elevation myocardial infarction and preserved left ventricular systolic function, comparison by two-dimensional speckle tracking versus tissue Doppler imaging of, 1146
 systolic, in patients with heart failure, relation to left ventricular ejection fraction and to QRS duration of, 602
 three-dimensional global mechanical, impact of right ventricular volume overload after surgical repair of tetralogy of Fallot on, 1731
 Left ventricular ejection fractions. *See* Ejection fractions, left ventricular
 Left ventricular filling pressure, predicting noninvasively, 371 (Readers' Comments)
 Left ventricular function. *See also* Left ventricular dysfunction
 after isolated aortic valve replacement for aortic valve stenosis, usefulness of left ventricular shape to predict the early recovery of, 1530
 diastolic, in trained male weightlifters at rest and during isometric exercise, 97
 effects of interruption of long-term cardiac resynchronization therapy on left ventricular dyssynchrony and, 718
 left ventricular noncompaction correlation with, new classification scheme for, 92
 myocardial velocities measured by two-dimensional speckle tracking versus tissue Doppler imaging of, 784
 operation for Ebstein anomaly and, 1724
 preserved systolic, in patients with non–ST-elevation myocardial infarction, comparison of left ventricular dyssynchrony by two-dimensional speckle tracking versus tissue Doppler imaging in, 1146
 quantitation by real-time three-dimensional echocardiography versus cardiac magnetic resonance, accuracy and reproducibility of, 778
 regional radial, relation between radial wall motion abnormalities using two-dimensional speckle tracking in children with idiopathic dilated cardiomyopathy and, 335
 systolic and diastolic, in asymptomatic patients with moderate aortic stenosis, 897
- Left ventricular geometry, progression of hypertensive heart disease association with myocardial strain and, 87
- Left ventricular hypertrophy, differentiation with three-dimensional echocardiography of hypertrophic cardiomyopathy from other forms of, 616
- Left ventricular mass
 index of, increased, in patients with stable coronary heart disease, prognostic significance to mortality and sudden death of (from the Heart and Soul Study), 1131
 left atrial size in National Football League linemen versus nonlinemen and, 343
- Left ventricular noncompaction, new classification scheme of ventricular performance and, 92
- Left ventricular volume, relation between mitral valve annulus and coronary sinus, influence of left atrial volume and, 890
- Left ventricular wall
 motion changes, invasively assessed microvascular integrity indexes in patients with ST-elevation myocardial infarction, predictive accuracy of fluorine-18 fluorodeoxyglucose positron emission tomography versus, 129
 right ventricular cardiomyocyte apoptosis in patients with acute myocardial infarction of, 658
 thickness, usefulness of changes in, prediction of full or partial pressure reperfusion in ST-elevation acute myocardial infarction and, 249
- Levine, Samuel A., history of grading systolic murmurs and, 1107
- Levosimendan, for men with acutely decompensated chronic heart failure, dobutamine-induced changes of two-dimensional longitudinal deformation usefulness to predict clinical and neurohumoral improvement in, 1225
- Lifestyle modifications
 comprehensive therapeutic, effect on prehypertension of, 1677
 socioeconomic status among survivors of acute myocardial infarction and, 1583
- Lipid-lowering therapy
 effectiveness of thrice weekly ezetimibe, 1205
 Effect of Combination Ezetimibe and High-Dose Simvastatin versus Simvastatin Alone on the Atherosclerotic Process in Patients With Heterozygous Familial Hypercholesterolemia (ENHANCE) trial, public hype and careful analysis of, 368 (Editorial)
 intensive, for patients with aortic stenosis, 1571

- in patients presenting with acute myocardial infarction, relation between infarct size (creatinine kinase-MB level) and, 1119
- Lipids, serum
 comparison in subjects with and without metabolic syndrome, 424
 management beyond statins—reducing residual cardiovascular risk, 559 (Editor's Roundtable)
 treating, in the general population, 372 (Readers' Comments)
- Lipoprotein(a), cardiorespiratory fitness in patients with type 2 diabetes mellitus and serum levels of, 700
- Long QT syndrome
 cardiac event risk in patients with asthma treated with beta₂ agonists and, 866
 methadone-induced, role of implantable cardioverter-defibrillators in patients with, 370 (Readers' Comments)
- Loop recorders, implantable, reproducibility of electrocardiographic findings in patients with suspected reflex neurally-mediated syncope using, 1518
- Lovastatin, pleiotropic effect of, with and without cholestyramine, in the Post Coronary Artery Bypass Graft (Post CABG) Trial, 1023
- Low-density lipoprotein cholesterol. *See* Cholesterol, low-density lipoprotein
- Lupus erythematosus, systemic, coronary artery calcium and C-reactive protein in asymptomatic women with rheumatoid arthritis or, 755
- Magnetic resonance imaging, cardiac
 accuracy and reproducibility of left ventricular function quantitation by real-time three-dimensional echocardiography versus, 778
 velocity-encoded, tissue Doppler imaging for measurement of myocardial velocities, assessment of left ventricular dyssynchrony, and estimation of left ventricular filling pressures in patients with ischemic cardiomyopathy versus, 1366
- Medical history
 Empress Sissi and cardiac tamponade, 1278
 Levine, Samuel A., grading systolic murmurs and, 1107
- Medicare Category III transaction codes, analysis of, costs and clinical outcomes of coronary multidetector computed tomographic angiography in individuals without coronary artery disease from, 672
- Men
 with acutely decompensated chronic heart failure, dobutamine-induced changes of two-dimensional longitudinal deformation usefulness to predict clinical and neurohumoral improvement after levosimendan treatment in, 1225
 age and double product (systolic blood pressure × heart rate) reserve-adjusted modification of the Duke Treadmill Score nomogram in, 1407
 alcohol consumption and heart failure in hypertensive United States physicians, 593
 apparently healthy physicians, change in high-density lipoprotein cholesterol and incident coronary heart disease in, 1663
 comparison of clinical features of arrhythmogenic right ventricular cardiomyopathy in women versus, 1252
 with congenital heart disease, erectile dysfunction in, 1728
 with coronary artery disease, peripheral arterial response compared to mental stress in women versus, 970
 defining normal distributions of coronary artery calcium in women and (from the Framingham Heart Study), 1136
 left ventricular diastolic function in trained weightlifters at rest and during isometric exercise, 97
 with severe high-density lipoprotein cholesterol deficiency, comparison of treatment with daily atorvastatin (20 mg) versus fenofibrate (200 mg) versus extended-release niacin (2 g) for, 1341
- Metabolic syndrome
 adverse effects on carotid artery intima-media thickness of, parental coronary artery disease and (from the Bogalusa Heart Study), 180
 all-cause and cardiovascular mortality relation to the number of risk factors due to, 689
 all-cause and cardiovascular mortality using the different definitions of, 188
 biomarkers of inflammation and oxidative stress with, effect of 10 mg versus 80 mg atorvastatin on, 321
 comparison of serum lipid values in subjects with and without, 424
 diagnostic usefulness of body mass index versus waist circumference in identifying apparently healthy subjects at increased risk of cardiovascular disease, 40
- Metalloproteinase-9, plasma activity in non-ST-segment elevation acute coronary syndrome patients with intensive atorvastatin therapy and, 12
- Metastasis, of hepatocellular carcinoma to heart, case report and review of the literature, 643
- Methadone, long QT syndrome induced by, role of implantable cardioverter-defibrillators in, 370 (Readers' Comments)
- Microalbuminuria, foscipril and serum C-reactive protein levels in patients with, 223
- Microvolt T-wave alternans
 comparison of the initial screening with repeat testing, long-term prognosis and, 1114 (Readers' Comments)
 in ischemic cardiomyopathy, usefulness to predict outcomes beyond one year of, 280
- Migraine headache, relief following, patent foramen ovale "closure" despite residual right-to-left shunt and, 916
- Mitral regurgitation
 asymptomatic or mildly symptomatic, exercise-induced functional changes in patients with idiopathic dilated cardiomyopathy of, 481
 severe, effect of beta-blocker therapy in patients with normal left ventricular ejection fraction and, 611
 valve replacement for aortic stenosis and, 1378
- Mitral valve annulus, coronary sinus and, influence of left atrial and ventricular volumes on, 890
- Modification of Diet in Renal Disease equation, mortality prediction in patients admitted for exclusion of myocardial ischemia with Cockcroft-Gault equation versus, 140
- Monitoring trends and determinants on Cardiovascular diseases (MONICA)/Cooperative Health Research in the Region of Augsburg (KORA) Augsburg Myocardial Infarction Registry, differential impact of admission C-reactive protein levels on 28-day mortality risk in patients with ST-elevation myocardial infarction versus non-ST-elevation myocardial infarction, 1125
- Mortality. *See also* Cardiac death, sudden
 28-day risk, differential impact of admission C-reactive protein levels in patients with ST-elevation versus non-ST-elevation myocardial infarction on (from the monitoring trends and determinants on cardiovascular diseases [MONICA)/Cooperative Health Research in the Region of Augsburg [KORA] Augsburg Myocardial Infarction Registry), 1125
 acute, in hospitalized patients undergoing echocardiography with and without an ultrasound contrast agent (Multicenter Registry Results in 4,300,966 Consecutive Patients), 1742
 after drug-eluting versus bare-metal stent coronary intervention in patients with diabetes mellitus, 165
 all-cause, in patients with recent cerebrovascular or coronary ischemic events, relation between aspirin dose, bleeding and (from the BRAVO Trial), 1285
 all-cause and cardiovascular, relation of the number of metabolic syndrome risk factors with, 689
 all-cause and cardiovascular, using the different definitions of metabolic syndrome, 188
 all-cause in-hospital and six-month, admission neutrophil to lymphocyte ratio in acute coronary syndrome patients and, 653
 blood transfusion in acute myocardial infarction and, 1116 (Readers' Comments)
 with Brugada-type electrocardiograms in a population of middle-aged Japanese diagnosed during a health examination, 584
 cardiac resynchronization therapy in patients with atrial fibrillation versus patients in sinus rhythm (Results of the Spanish Atrial Fibrillation and Resynchronization [SPARE] Study), 444
 cardiovascular, in patients older than 65 years, chronic kidney disease, diabetes mellitus, and myocardial infarction as equivalent risk factors for, 1668
 cardiovascular, paradoxical systolic blood pressure increase after exercise as a predictor of, 518
 comparison of the impact of short (<1 year) and long-term (≥1 year) clopidogrel use following percutaneous coronary intervention on, 1159
 failure to prescribe beta blockers acutely in patients with sustained ventricular tachycardia and ventricular fibrillation following acute myocardial infarction and (from the VALsartan In Acute myocardial infarction trial [VALIANT] Registry), 1427

- fasting plasma glucose usefulness to predict coronary heart disease in persons ≥ 60 years of age without diabetes mellitus or in those with undiagnosed diabetes mellitus (from The Dubbo Study), 825
- in hospitalized heart failure patients with left ventricular ejection fraction $>50\%$ versus those with ejection fraction $<50\%$, 79
- in-hospital, troponin I levels in patients with ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage and, 632
- in-hospital and late postoperative, in patients undergoing aortic valve replacement for symptomatic aortic stenosis, comparison of brain natriuretic peptide versus logistic EuroSCORE in predicting, 749
- in-hospital and long-term, impact of early and delayed stroke after isolated coronary artery bypass grafting on, 411
- late, in hospitalized patients >70 years of age with systolic heart failure receiving versus not receiving beta blockers, 1711
- left atrial volume index versus left ventricular ejection fraction usefulness to predict heart failure hospitalization in ambulatory patients with coronary heart disease and (from the Heart and Soul Study), 70
- in patients with implantable cardioverter-defibrillators, effect of beta blockers, angiotensin-converting enzyme inhibitors or angiotensin receptor blockers, and statins on, 77
- in patients with intermediate to high risk of coronary artery disease, relation of exercise capacity and body mass index to, 1028
- in patients with peripheral arterial disease with versus without associated chronic obstructive pulmonary disease, statin therapy effect on, 192
- prediction in ischemic cardiomyopathy beyond one year using microvolt T-wave alternans, 280
- prediction with Modification of Diet in Renal Disease versus Cockcroft-Gault equations in patients admitted for exclusion of myocardial ischemia, 140
- preoperative statins and outcomes in isolated cardiac valve surgery and, 1235
- prognostic significance of increased left ventricular mass index in patients with stable coronary heart disease to sudden death and (from the Heart and Soul Study), 1131
- proteinuria usefulness as prognostic marker among patients undergoing percutaneous coronary intervention of (data from the Evaluation of Oral Xemilofiban in Controlling Thrombotic Events [EXCITE] Trial), 1151
- serum uric acid levels in patients with suspected or definite coronary artery disease and, 797
- troponin T elevation in patients without chest pain and, 668
- Mueller maneuver, sleep apnea, atrial fibrillation, and heart failure implications from dynamic changes of left ventricular performance and left atrial volume induced in healthy young adults using, 1557
- Multicenter Registry Results in 4,300,966 Consecutive Patients, acute mortality in hospitalized patients undergoing echocardiography with and without an ultrasound contrast agent, 1742
- Multidetector computed tomography. *See* Computed tomography
- Multi-Ethnic Study of Atherosclerosis (MESA), aortic wall thickness and distensibility relation to cardiovascular risk factors, 491
- Murmurs, systolic, Samuel A. Levine and the history of grading, 1107
- Myocardial bridging, of left anterior descending coronary artery, comparison of angiographic versus intravascular ultrasonic detection of, 1608
- Myocardial damage, frequency of elevation of cardiomyocyte damage markers in otherwise healthy newborns and, 761
- Myocardial infarction
- accuracy of coding now that coding descriptions have been clarified to distinguish ST-elevation myocardial infarction from non-ST elevation myocardial infarction, 513
 - acute (*See* Acute myocardial infarction)
 - after drug-eluting versus bare-metal stent coronary intervention in patients with diabetes mellitus, 165
 - cardiovascular mortality in patients older than 65 years and, 1668
 - cocaine use and, 966
 - non-ST-elevation
 - accuracy of coding myocardial infarction now that coding descriptions have been clarified to distinguish ST-elevation myocardial infarction from, 513
 - differential impact of admission C-reactive protein levels on 28-day mortality risk in patients with ST-elevation myocardial infarction versus (from the monitoring trends and determinants on cardiovascular diseases [MONICA]/Cooperative Health Research in the Region of Augsburg [KORA] Augsburg Myocardial Infarction Registry), 1125
 - hemorheological variables in ST-elevation myocardial infarction or unstable angina pectoris versus, 125
 - and preserved left ventricular systolic function, comparison of left ventricular dyssynchrony by two-dimensional speckle tracking versus tissue Doppler imaging in, 1146
 - perioperative, impact on angiographic and clinical outcomes following coronary artery bypass grafting of (from PProject of Ex-vivo Vein graft ENgineering via Transection [PREVENT] IV), 546
 - ST-segment elevation (*See* ST-segment elevation myocardial infarction)
- Myocardial ischemia
- by dobutamine stress echocardiography in patients without angina pectoris after coronary revascularization, prognostic significance of, 1156
 - mortality prediction with Cockcroft-Gault equation versus Modification of Diet in Renal Disease equations in patients admitted for exclusion of, 140
- Myocardial perfusion, at cardiac catheterization, quantitative automated assessment of, 980
- Myocardial perfusion imaging
- evaluation of patients for cardiac resynchronization therapy and, 211
 - in patients with end-stage renal disease undergoing coronary angiography, 1451
 - stress, effectiveness of anemia and chronic kidney disease as predictors for presence and severity of coronary artery disease in patients undergoing, 266
- Myocardial strain, progression of hypertensive heart disease association with left ventricular geometry and, 87
- National Football League, left ventricular mass and left atrial size in linemen versus nonlinemen of, 343
- National Health and Nutrition Examination Survey (NHANES) 2001 to 2004, hypovitaminosis D prevalence in cardiovascular diseases, 1540
- National Health and Nutrition Examination Survey (NHANES) I and II Studies, comparison of cardiovascular risk factors for high brachial pulse pressure in blacks versus whites, 1514
- National Heart, Lung, and Blood Institute Sponsored WISE Study, diabetes mellitus, hypothalamic hypoestrogenemia, and coronary artery disease in premenopausal women, 150
- Natriuretic peptides
- association between renal function and circulating levels of (from the Dallas Heart Study), 1394
 - B-type (*See also* N-terminal pro-B-type natriuretic peptide)
 - serum levels compared to logistic EuroSCORE in predicting in-hospital and late postoperative mortality in patients undergoing aortic valve replacement for symptomatic aortic stenosis, 749
 - tissue Doppler imaging for differentiation of constrictive pericardial disease from restrictive cardiomyopathy versus, 357
- Neoplasms, fatal. *See also* Carcinoid heart disease
- long-term risk following radiofrequency ablation in children of, 913
- Nephropathy, contrast-induced
- after cardiac catheterization, gender and age factors in, 1482
 - incidence in patients with chronic renal insufficiency undergoing multidetector computed tomographic angiography treated with preventive measures, 353
 - timing of testing renal function to disclose, 371 (Readers' Comments)
- Neutrophil to lymphocyte ratio, on admission, outcomes in acute coronary syndrome patients and, 653
- Niacin, extended-release (2 g), for severe high-density lipoprotein cholesterol deficiency in men, comparison of treatment with daily atorvastatin (20 mg) versus fenofibrate (200 mg) versus, 1341
- Nonsteroidal antiinflammatory drugs, cardiovascular risk and, 1046 (Editor's Roundtable)
- N-terminal pro-B-type natriuretic peptide. *See also* Natriuretic peptides, B-type
- prognostic usefulness in patients with heart failure with versus without chronic kidney disease, 469
 - tissue Doppler variables of left ventricle diastolic function and, prognostic value in patients with chronic systolic heart failure of, 463
 - usefulness as biomarker for carcinoid heart disease, 938

- usefulness of brain natriuretic peptide to predict cardiovascular outcomes in patients with heart failure and preserved left ventricular ejection fractions and, 733
- Obese patients**
 bariatric surgery for, cardiovascular risk after, 930
 morbidly, body weight limitations of United States cardiac catheterization laboratories for, 285
- Occluders, plug, transcatheter closure of patent ductus arteriosus in children using the swivel-disk device versus, 486**
- Omega-3 acid ethyl esters, prescription addition to simvastatin, effect on lipids and lipoprotein particles in men and women with mixed dyslipidemia and, 429, 1425 (Correction)**
- Omega-3 fatty acids, for mixed dyslipidemia, combined statin therapy plus, 1040**
- Ongoing Telmisartan Alone and in Combination With Ramipril Global Endpoint Trial (ONTARGET), angiotensin receptor blockers and cardiovascular protection, 1281 (Editorial)**
- Optical coherence tomography**
 assessment of culprit and remote coronary narrowings with long-term outcomes using, 391
 distribution and frequency of thin-capped fibroatheromas and ruptured plaques in the entire culprit coronary artery in patients with acute coronary syndrome as determined by, 975
 neointimal coverage comparison of sirolimus-eluting stent versus bare-metal stent three months after implantation, 27
- Organized Program to Initiate Lifesaving Treatment in Hospitalized Patients With Heart Failure, beta-blocker therapy dosing before, during, and after hospitalization for heart failure, 1524**
- Outcome assessment, after cardiac surgery, continuous quality improvement program and, 772**
- Oxygen uptake, peak, reproducibility in heart failure patients (from the Heart Failure and A Controlled Trial Investigating Outcomes of exercise training), 712**
- Pacemakers**
 dual-chamber, implantation in patients without a previous history of atrial fibrillation, frequency, duration, and predictors of newly-diagnosed atrial fibrillation in, 450
 lead for, management of subacute and delayed right ventricular perforation with, 1352
- Paclitaxel-eluting stents**
 clinical and angiographic follow-up of small vessel lesions treated with (from the TRUE Registry), 1002
 impact of mild or moderate renal insufficiency on intravascular ultrasound analysis of chronic vascular response to bare-metal stents and (from the TAXUS IV, V, and VI Trials), 1009
 Taxus Express-2, intravascular ultrasound assessment of expansion of sirolimus-eluting stent (Cypher Select) in patients with diabetes mellitus versus, 19
 two-year outcome of patients treated with sirolimus-eluting stents in an unselected population with coronary artery disease versus (from the REWARDS Registry), 292
- Patent ductus arteriosus, transcatheter closure using the swivel-disk device versus plug occluder in children, 486**
- Patent foramen ovale, closure, migraine headache relief despite residual right-to-left shunt after, 916**
- Pediatric cardiology. See Children**
- Percutaneous coronary intervention(s) (PCI). See also Saphenous vein grafts**
 for acute coronary syndrome, clopidogrel benefit doubt due to rebound adverse events after, 248 (Readers' Comments)
 for acute coronary syndrome in chronic kidney disease patients on dialysis, impact of optimal medical therapy on outcome of, 535
 clopidogrel loading dose in healthy volunteers, and platelet functional recovery dynamics after, 790
 comparison of myocardial reperfusion in patients with fasting blood glucose ≤ 100 , 101 to 125, and > 125 mg/dl and ST-elevation myocardial infarction with, 1457
 drug-eluting stent safety in patients with left ventricular dysfunction undergoing, 679
 with drug-eluting versus bare-metal stents in diabetes mellitus, stent thrombosis, myocardial infarction and mortality after, 165
 impact of 600-mg loading dose of clopidogrel on 30-day outcome in unselected patients undergoing, 1318
 long-term outcomes of bifurcation narrowings after, 404
 mortality after, comparison of the impact of short (< 1 year) and long-term (≥ 1 year) clopidogrel on, 1159
 offsite, safety and efficacy in 1,348 consecutive patients in rural Tasmania, 1323
 previous, comparison of effectiveness of atorvastatin 10 mg versus 80 mg in reducing major cardiovascular events and repeat revascularization in patients with (Post Hoc Analysis of the Treating to New Targets [TNT Study]), 1312
 primary
 for acute ST-elevation myocardial infarction, patterns of upstream antiplatelet therapy use before (from the CRUSADE National Quality Improvement Initiative), 1335
 clinical and electrocardiographic correlates of normal coronary angiography in patients referred for, 155
 off-site, Dutch randomized trial on, 1115 (Readers' Comments)
 for ST-elevation acute myocardial infarction, predictive value comparison of four different risk scores for outcomes in, 6
 as primary revascularization in patients with acute coronary syndrome, coronary artery bypass grafting versus, 173
 proteinuria usefulness as a prognostic marker of mortality and cardiovascular events among patients undergoing (data from the Evaluation of Oral Xemilofiban in Controlling Thrombotic Events [EXCITE] Trial), 1151
 repeated, angiographic patterns of drug-eluting stent restenosis and one-year outcomes after treatment with, 311
 risk predictors of retroperitoneal hemorrhage following, 1473
 same-day, disparities in use for patients with ST-elevation myocardial infarction in Florida, 2001–2005, 802
 for stable or unstable angina pectoris, bleeding risk and outcomes of bivalirudin versus glycoprotein IIb/IIIa inhibitors with targeted low-dose unfractionated heparin in, 160
 for ST-elevation myocardial infarction, emergency pretreatment for contrast allergy before, 1469
 stress echocardiography usefulness for risk stratification of patients after, 1170
 suboptimal, 954 (Readers' Comments)
 successful versus unsuccessful, comparison of safety, efficacy, and outcome in "true" chronic total occlusions of, 1175
 time from symptom onset to balloon inflation by, relation of left ventricular infarct transmural and infarct size to, 1163
 using drug-eluting stents, comparison of higher clopidogrel loading and maintenance dose to standard dose on platelet function and outcomes after, 401
- Periodontal disease, self-reported, usefulness to identify individuals with elevated inflammatory markers at risk of cardiovascular disease, 1509**
- Peripheral arterial disease**
 physical activity during daily life and circulating biomarker levels in patients with, 1263
 usefulness of hypertensive blood pressure response during single-stage exercise testing to predict long-term outcome in patients with, 921
 with versus without associated chronic obstructive pulmonary disease, statin therapy effect on mortality in patients with, 192
- Peripheral arterial vasoconstriction, mental stress in women versus men with coronary artery disease compared to, 970**
- Peroxisome proliferator activated receptor alpha agonist, potent and selective, efficacy and safety in subjects with type 2 diabetes mellitus and dyslipidemia, 434**
- Physical activity. See also Exercise**
 during daily life and circulating biomarker levels in patients with peripheral arterial disease, 1263
 in patients with systemic right ventricle, cardiac function, exercise capacity, quality of life and, 1258
 self-reported leisure-time, usefulness to long-term survival prediction in patients with coronary heart disease, 375
- Physicians**
 apparently healthy male, change in high-density lipoprotein cholesterol and incident coronary heart disease in, 1663

- estimation of patient cardiovascular risk by, discordance between evidence-based medical therapy and, 1142
- time to diagnosis for acute aortic dissection by, 1399
- Plaques**
- noncalcified coronary, comparison of 64-slice multidetector computed tomography and spectral analysis of intravascular ultrasound backscatter signals of, 988
 - ruptured, in patients with acute coronary syndrome, distribution and frequency in the entire culprit coronary artery as determined by optical coherence tomography of, 975
 - vulnerable, 1644 (Editor's Roundtable)
- Platelet aggregation, adenosine diphosphate-induced, reloading effectiveness to overcome clopidogrel nonresponsiveness in patients with acute myocardial infarction and, 524**
- Platelet functional recovery dynamics, clopidogrel loading dose in healthy volunteers and, 790**
- Plug occluder, transcatheter closure of patent ductus arteriosus in children using the swivel-disk device versus, 486**
- Positron emission tomography, fluorine-18 fluorodeoxyglucose, predictive accuracy of left ventricular wall motion changes in patients with ST-elevation myocardial infarction of invasively assessed microvascular integrity indexes versus, 129**
- Post Coronary Artery Bypass Graft (Post CABG) Trial, pleiotropic effect of lovastatin, with and without cholestyramine, 1023**
- Post Hoc Analysis of the Treating to New Targets (TNT) Study, comparison of effectiveness of atorvastatin 10 mg versus 80 mg in reducing major cardiovascular events and repeat revascularization in patients with previous percutaneous coronary intervention, 1312**
- Poststent Optimal Stent Expansion Trial (POET)**
- correct optimization of drug-eluting stents and, 1113 (Readers' Comments)
 - intravascular ultrasound evaluation of optimal drug-eluting stent expansion after poststent balloon dilation using a noncompliant balloon versus a semi-compliant balloon, 304
- Poverty, and cardiovascular disease in India, 102**
- Pravastatin or Atorvastatin Evaluation and Infection Therapy (PROVE IT-TIMI 22), outcomes of patients with previous coronary artery bypass grafting, 552**
- Pregnancy-associated plasma protein A, as a quick and sensitive biomarker in early phase of acute coronary syndrome, 954 (Readers' Comments)**
- Prehospital delay, trends in patients with acute myocardial infarction (from the Worcester Heart Attack Study), 1589**
- Prehypertension. *See also* Hypertension**
- effect of comprehensive therapeutic lifestyle changes on, 1677
- Primary Angioplasty in Myocardial Infarction (PAMI), predictive value of, in primary percutaneous coronary intervention for outcomes in ST-elevation acute myocardial infarction comparison with three different risk scores, 6**
- Project of Ex-vivo Vein graft ENgineering via Transfection (PREVENT) IV, perioperative myocardial infarction impact on angiographic and clinical outcomes following coronary artery bypass grafting, 546**
- Prostaglandin E2, plasma levels in non-ST-segment elevation acute coronary syndrome patients with intensive atorvastatin therapy and, 12**
- Proteinuria, usefulness as a prognostic marker of mortality and cardiovascular events among patients undergoing percutaneous coronary intervention (data from the Evaluation of Oral Xemilofiban in Controlling Thrombotic Events [EXCITE] Trial), 1151**
- Proximal isovelocity surface area, of aortic regurgitation volume, with central versus eccentric jets, quantification accuracy using, 475**
- Psoriasis**
- coronary artery disease and, 1631 (Editor's Consensus)
 - inflammation, atrial fibrillation and, 647 (Readers' Comments)
- Pulmonary arterial hypertension**
- evidence for early right ventricular and septal mechanical activation (interventricular dyssynchrony) in, 1273
 - severe, complicating "isolated" atrial septal defect in infancy, 340
- P-wave dispersion in standard twelve-lead electrocardiography, prediction of transition from paroxysmal to persistent atrial fibrillation using, 573**
- QRS duration, in patients with heart failure, left ventricular systolic dyssynchrony relation to left ventricular ejection fractions and, 602**
- Quality of care, for heart failure, hospital length of stay and, 1693**
- Quality of life, in patients with systemic right ventricle, physical activity, cardiac function, exercise capacity and, 1258**
- Radiofrequency ablation, in children, long-term risk of fatal malignancy following, 913**
- Red blood cells, transfusion impact on clinical outcomes with acute myocardial infarction, 115**
- REGIMEN Registry, perioperative bridging therapy with unfractionated heparin or low-molecular-weight heparin in mechanical prosthetic valve patients on long-term oral anticoagulants, 879**
- Remote-controlled magnetic navigation, for ablation of atrial fibrillation, feasibility and safety of, 1674**
- Remote ischemic preconditioning, in cardiovascular surgery, review and meta-analysis of randomized controlled trials of, 1487**
- Renal disease, end-stage, myocardial perfusion imaging in patients undergoing coronary angiography with, 1451**
- Renal dysfunction, in chronic heart failure, telomere length and, 207**
- Renal function**
- association between circulating levels of natriuretic peptides and (from the Dallas Heart Study), 1394
 - timing of testing, to disclose contrast-induced nephropathy, 371 (Readers' Comments)
- Renal insufficiency**
- chronic
 - and contrast-induced nephropathy incidence in patients undergoing multidetector computed tomographic angiography treated with preventive measures, 353
 - retroperitoneal hemorrhage following percutaneous coronary intervention and, 1473
 - mild or moderate, impact on the intravascular ultrasonic analysis of chronic vascular response to paclitaxel-eluting and bare-metal stents (from the TAXUS IV, V, and VI Trials), 1009
- Reperfusion, full or partial pressure, in ST-segment elevation acute myocardial infarction, left ventricular wall thickness changes and prediction of, 249**
- Respiratory artifact, in electrocardiograms, significance of, 1090**
- Restenosis, drug-eluting stent, angiographic patterns after treatment with repeated percutaneous coronary intervention of one-year outcomes and, 311**
- Resynchronization therapy. *See* Cardiac resynchronization therapy**
- Retinal vascular caliber, coronary heart disease risk prediction and (from the Atherosclerosis Risk In Communities [ARIC] Study), 58**
- REWARDS Registry, two-year outcome of patients treated with sirolimus- versus paclitaxel-eluting stents in an unselected population with coronary artery disease, 292**
- Rheumatoid arthritis, coronary artery calcium and C-reactive protein in asymptomatic women with systematic lupus erythematosus or, 755**
- Right ventricle**
- acute myocardial infarction of left ventricular wall and cardiomyocyte apoptosis of, 658
 - new generation of three-dimensional echocardiography for volumetric and functional measurements of, 499
 - subacute and delayed perforation with a pacing or implantable cardioverter-defibrillator lead, management of, 1352
 - systemic, relation of physical activity, cardiac function, exercise capacity, and quality of life in patients with, 1258
- Right ventricular volume, overload impact on three-dimensional global left ventricular mechanical dyssynchrony after surgical repair of tetralogy of Fallot, 1731**
- Rosuvastatin, meta-analysis of head-to-head randomized controlled trials of benefits and risks of atorvastatin versus, 1654**
- Roundtable discussions, informativeness and usefulness of, 507 (Readers' Comments)**
- Runners, long-distance, aged ≥ 55 years, magnitude, reproducibility, and association with baseline cardiac function of cardiac biomarker release in, 218**

- Saphenous vein grafts. *See also* Percutaneous coronary intervention(s) drug-eluting stents versus bare-metal stents for narrowing in, 530
- Sclerosis, stage-related effect of statin treatment on the progression of aortic valve stenosis and, 738
- Seasonal variations, incidence and case fatality of acute myocardial infarction in a Japanese Population (from the Takashima AMI Registry, 1988 to 2003), 1307
- Sedentary occupations, endothelial cell function predictors in employees in a worksite exercise program and, 820
- Septal defect. *See* Atrial septal defect; Ventricular septal defect
- Shock. *See* Cardiogenic shock
- Simvastatin
 - atherosclerotic process in patients with heterozygous familial hypercholesterolemia and (from the ENHANCE trial), public hype and careful analysis of, 368 (Editorial)
 - lipids and lipoprotein particles in men and women with mixed dyslipidemia and addition of prescription omega-3 acid ethyl esters to, 429, 1425 (Correction)
- Sinus rhythm
 - real-time three-dimensional echocardiography comparison of left atrial volumes and function in patients having catheter ablation for atrial fibrillation with recurrent atrial fibrillation three months later versus persistence of, 847
 - restoration in sleep apnea patients with atrial fibrillation or flutter and, 709
- Sirolimus-eluting stents
 - (Cypher Select), intravascular ultrasound assessment of expansion of paclitaxel-eluting (Taxus Express-2) stent in patients with diabetes mellitus versus, 19
 - three months after implantation, neointimal coverage by optical coherence tomography of bare-metal stents versus, 27
 - two-year outcome of patients treated with paclitaxel-eluting stents in an unselected population with coronary artery disease versus (from the REWARDS Registry), 292
- Sleep apnea
 - dynamic changes of left ventricular performance and left atrial volume induced in healthy young adults using the Mueller maneuver and implications for, 1557
 - sinus rhythm restoration in patients with atrial fibrillation or flutter and effect on, 709
- Sleep-disordered breathing
 - life-threatening ventricular arrhythmia in heart failure patients with implantable cardioverter-defibrillator and, 1064
- Smoking, cardiovascular effect of bans in public places on, 1421 (Editorial)
- Socioeconomic status, lifestyle modifications among survivors of acute myocardial infarction and, 1583
- Sodium, dietary
 - hospital readmission in patients with heart failure, and lack of knowledge about, 1212
- Soy nuts
 - adhesion molecules and inflammation markers in hypertensive and normotensive postmenopausal women and, 84
- Spanish Atrial Fibrillation and Resynchronization (SPARE) Study
 - benefits and mortality of cardiac resynchronization therapy in patients with atrial fibrillation versus patients in sinus rhythm, 444
- Speckle tracking, two-dimensional
 - in children with idiopathic dilated cardiomyopathy to assess left ventricular regional radial function relation to radial wall motion abnormalities, 335
 - myocardial tissue velocities measured by tissue Doppler imaging versus, 784
 - in patients with non-ST-elevation myocardial infarction and preserved left ventricular systolic function, comparison of left ventricular dyssynchrony by tissue Doppler imaging versus, 1146
- Statins. *See also* Atorvastatin; Lovastatin; Simvastatin
 - aortic valve sclerosis and stenosis, progression of, stage-related effect of treatment with, 738
 - intensive, for patients with aortic stenosis, 1571
 - lipid management beyond, reducing cardiovascular residual risk and, 559 (Editor's Roundtable)
 - for mixed dyslipidemia, combined omega-3 fatty acid therapy plus, 1040
 - mortality in patients with implantable cardioverter-defibrillators and, 77
 - mortality in patients with peripheral arterial disease, with versus without associated chronic obstructive pulmonary disease, 192
 - preoperative, outcomes in patients undergoing isolated cardiac valve surgery and, 1235
 - risk of heart failure after acute myocardial infarction and, 1706
- STEMI. *See* ST-segment elevation myocardial infarction
- Stenosis. *See also* Restenosis
 - aortic
 - intensive lipid-lowering therapy for patients with, 1571
 - mitral regurgitation and valve replacement for, 1378
 - usefulness of left ventricular shape to predict early recovery of left ventricular function after isolated aortic valve replacement for, 1530
 - asymptomatic aortic, atorvastatin modulation of prognosis and risk factors in patients with, 743
 - moderate aortic, left ventricular systolic and diastolic function in asymptomatic patients with, 897
 - severe aortic
 - feasibility and initial results of percutaneous aortic valve implantation including selection of transfemoral or transapical approach in patients with, 1240
 - stage-related effect of statin treatment on the progression of aortic valve sclerosis and, 738
 - unprotected left main coronary artery, two-year clinical outcome with drug-eluting stents versus bare-metal stents in a real-world registry of (from the Italian Society of Invasive Cardiology), 1463
- Stents
 - bare metal (*See* Bare-metal stents)
 - bifurcation, long-term outcomes of narrowings after percutaneous coronary intervention, 404
 - drug-eluting
 - bare metal stents for narrowing in saphenous vein grafts versus, 530
 - comparison of higher clopidogrel loading and maintenance dose to standard dose on platelet function and outcomes after percutaneous coronary intervention using, 401
 - correct optimization of, Poststent Optimal Stent Expansion Trial (POET) and, 1113 (Readers' Comments)
 - effect of prolonged thienopyridine use after implantation of (from the TAXUS Landmark Trials Data), 1017
 - implantation
 - incidence of bleeding and compliance on prolonged dual antiplatelet therapy (aspirin + thienopyridine) following, 1477
 - "nuisance" bleeding and clopidogrel compliance in patients undergoing, 1614
 - indications and complications of bare-metal stents versus, 32 (Editor's Roundtable)
 - intravascular ultrasound evaluation of optimal expansion after poststent balloon dilation using a noncompliant balloon versus a semicompliant balloon (from the Poststent Optimal Stent Expansion Trial [POET]), 304
 - newly placed, virtual histology intravascular ultrasound appearance of, 1182
 - in patients with diabetes mellitus, meta-analysis comparison (nine trials) of outcomes with bare-metal stents versus, 1328
 - restenosis and one-year outcomes after treatment with repeated percutaneous coronary intervention, angiographic patterns of, 311
 - safety in patients with left ventricular dysfunction undergoing percutaneous coronary intervention, 679
 - thrombosis, definite and probable, at a single hospital in a four-year period, observations and outcomes of, 298
 - thrombosis, myocardial infarction, and mortality after bare-metal stent coronary intervention in patients with diabetes mellitus versus, 165
 - thrombosis incidence, based on Academic Research Consortium definitions, 683
 - two-year clinical outcome with bare-metal stents in a real-world registry of unprotected left main coronary artery stenosis versus (from the Italian Society of Invasive Cardiology), 1463
 - everolimus-eluting, Xience V, overview of the 2007 Food and Drug Administration Circulatory System Devices Panel Meeting on, 1624
 - implantation of, long-term outcomes in patients on dual oral antiplatelet treatment requiring oral anticoagulant therapy, 1618

- paclitaxel-eluting (*See* Paclitaxel-eluting stents)
- prognostic value of lesion classification for the clinical outcome after placement of, 954 (Readers' Comments)
- pulmonary artery, hybrid intraoperative, for congenital heart disease, 1737
- sirolimus-eluting (*See* Sirolimus-eluting stents)
- thrombosis prediction using a clinical risk score, 541
- Stimulants, illicit
- use in a United States heart failure population presenting to the emergency department (from the Acute Decompensated Heart Failure National Registry Emergency Module), 1216
- Stress
- mental, peripheral arterial response in women versus men with coronary artery disease compared to, 970
- oxidative, oxidized low-density lipoproteins and, 1118 (Readers' Comments)
- Stress testing. *See* Exercise testing, stress
- Stroke
- adult-onset asthma relation to coronary heart disease and, 508 (Readers' Comments)
- early and delayed, impact on in-hospital and long-term mortality after isolated coronary artery bypass grafting, 411
- ischemic
- echocardiographic and clinical risk factors for atrial fibrillation in hypertensive patients with, 1348
- relation of troponin I levels to in-hospital mortality in patients with intracerebral hemorrhage, subarachnoid hemorrhage, and, 632
- ST-segment elevation myocardial infarction (STEMI). *See also* Acute myocardial infarction
- accuracy of coding myocardial infarction now that coding descriptions have been clarified to distinguish non-ST-elevation myocardial infarction from, 513
- acute
- changes in left ventricular wall thickness to predict full or partial pressure reperfusion in, 249
- undergoing primary percutaneous coronary intervention, predictive value comparison of four different risk scores for outcomes of patients with, 6
- acute, patterns of upstream antiplatelet therapy use before primary percutaneous coronary intervention for (from the CRUSADE National Quality Improvement Initiative), 1335
- body mass index and waist-to-hip ratio impact on clinical outcomes in patients with (from the Korean Acute Myocardial Infarction Registry), 957
- comparison of predictive accuracy of left ventricular wall motion changes between invasively assessed microvascular integrity indexes and fluorine-18 fluorodeoxyglucose positron emission tomography in patients with, 129
- differential impact of admission C-reactive protein levels on 28-day mortality risk in patients with non-ST-elevation myocardial infarction versus (from the monitoring trends and determinants on cardiovascular diseases [MONICA]/Cooperative Health Research in the Region of Augsburg [KORA] Augsburg Myocardial Infarction Registry), 1125
- disparities in same-day percutaneous coronary intervention for patients in Florida, 2001–2005, 802
- emergency pretreatment for contrast allergy before direct percutaneous coronary intervention for, 1469
- hemorheological variables in non-ST-elevation myocardial infarction or unstable angina pectoris versus, 125
- percutaneous coronary intervention for, comparison of myocardial reperfusion in patients with fasting blood glucose ≤ 100 , 101 to 125, and >125 mg/dl, 1457
- relation of left ventricular infarct transmural and infarct size after primary percutaneous coronary angioplasty to time from symptom onset to balloon inflation in, 1163
- Surgery. *See also* Vascular surgery
- bariatric, for obesity, cardiovascular risk after, 930
- cardiac, continuous quality improvement program and major morbidity after, 772
- cardiovascular, review and meta-analysis of randomized controlled clinical trials of remote ischemic preconditioning in, 1487
- coronary, dofetilide as routine prophylaxis for postoperative atrial fibrillation prevention after, 506 (Readers' Comments)
- isolated cardiac valve, preoperative statins and outcomes in, 1235
- relationship of inferior vena cava filter usage by various surgical and medical subspecialists after, 226
- Surgical revascularization. *See* Coronary artery bypass grafting
- Survival
- in both ischemic and nonischemic systolic heart failure, systolic blood pressure and, 1698
- long-term prediction in patients with coronary heart disease, self-reported leisure-time physical activity usefulness to, 375
- Swivel-disk device
- transcatheter closure of patent ductus arteriosus in children using the plug occluder versus, 486
- Syncope
- reproducibility of electrocardiographic findings in patients with suspected reflex neurally-mediated, 1518
- Takashima AMI Registry, 1988 to 2003, seasonal pattern of incidence and case fatality of acute myocardial infarction in a Japanese population, 1307
- TAXUS IV, V, and VI Trials, impact of mild or moderate renal insufficiency on intravascular ultrasound analysis of chronic vascular response to paclitaxel- and bare-metal stents, 1009
- TAXUS Landmark Trials Data, effect of prolonged thienopyridine use after drug-eluting stent implantation, 1017
- Telmisartan, cardiovascular protection and, from the Ongoing Telmisartan Alone and in Combination With Ramipril Global Endpoint Trial (ONTARGET), 1281 (Editorial)
- Telomere length, in chronic heart failure, renal dysfunction and, 207
- Tendon xanthoma, in clinical diagnosis for familial hypercholesterolemia, genetic testing versus, 1187
- Tetralogy of Fallot
- repaired, in adolescents, autonomic heart rate control at rest and during unloading of the right ventricle in, 1085
- surgical repair of, right ventricular volume overload on three-dimensional global left ventricular mechanical dyssynchrony after, 1731
- TGFBR1 and TGFBR2 mutations, absence of, in patients with bicuspid aortic valve and aortic dilation, 629
- Thalassemia intermedia, relation between iron-overload indices, cardiac echodoppler, and biochemical markers in, 363
- Thienopyridine
- and aspirin following drug-eluting stent implantation, incidence of bleeding and compliance with, 1477
- prolonged use of, after drug-eluting stent implantation (from the TAXUS Landmark Trials Data), 1017
- Thrombolysis In Myocardial Infarction (TIMI) score, predictive value of, in primary percutaneous coronary intervention for outcomes in ST-elevation acute myocardial infarction comparison with three different risk scores, 6
- Thrombosis
- stent
- after drug-eluting versus bare-metal stent coronary intervention in patients with diabetes mellitus, 165
- clinical risk score for prediction of, 541
- drug-eluting
- definite and probable at a single hospital in a four-year period, observations and outcomes of, 298
- incidence of, based on Academic Research Consortium definitions, 683
- vena cava, incidence in the United States, 927
- Tissue Doppler imaging. *See also* Echocardiography
- brain natriuretic peptide for differentiation of constrictive pericardial disease from restrictive cardiomyopathy versus, 357
- left ventricle diastolic function and N-terminal pro-type-B natriuretic peptide variables, prognostic value in patients with chronic systolic heart failure of, 463
- myocardial tissue velocities measured by two-dimensional speckle tracking versus, 784
- in patients with non-ST-elevation myocardial infarction and preserved left ventricular systolic function, comparison of left ventricular dyssynchrony by two-dimensional speckle tracking versus, 1146
- prognostic value of pulse-wave parameters in systolic heart failure, 722

- in thalassemia intermedia, relation between iron-overload indices, biochemical markers and, 363
- velocity-encoded magnetic resonance imaging for measurement of myocardial velocities, assessment of left ventricular dyssynchrony, and estimation of left ventricular filling pressures in patients with ischemic cardiomyopathy versus, 1366
- Transesophageal echocardiography, cardioversion guided by, anticoagulant and anti-inflammatory responses of enoxaparin versus unfractionated heparin for atrial fibrillation in, 842
- Troponin, prognostic implications of normal (<0.10 ng/ml) and borderline (0.10 to 1.49 ng/ml) elevation levels in critically ill patients without acute coronary syndrome of, 509
- Troponin I
 - clinical severity, cardiac dysfunction, and pulmonary congestion in patients with subarachnoid hemorrhage and, 1545
 - serum levels of, relation with in-hospital mortality in patients with ischemic stroke, intracerebral hemorrhage, and subarachnoid hemorrhage, 632
- Troponin T, elevation in patients without chest pain, prognostic significance of, 668
- TRUE Registry, clinical and angiographic follow-up of small vessel lesions treated with paclitaxel-eluting stents, 1002
- Ultrasound, intravascular. *See also* Echocardiography
 - analysis of chronic vascular response to paclitaxel-eluting and bare-metal stents, impact of mild or moderate renal insufficiency on (from the TAXUS IV, V, and VI Trials), 1009
 - angiographic detection of myocardial bridging of the left anterior descending coronary artery compared to, 1608
 - to assess the significance of angiographically ambiguous coronary narrowings, comparison of contrast-enhanced 64-slice computed tomography to, 994
 - contrast agents, safety in stress echocardiography, 1269
 - of optimal drug-eluting stent expansion after poststent balloon dilation using a noncompliant balloon versus a semicompliant balloon (from the Poststent Optimal Stent Expansion Trial [POETI]), 304
 - spectral analysis of backscatter signals of noncalcified coronary plaques, comparison of 64-slice multidetector computed tomography to, 988
 - virtual histology appearance of newly placed drug-eluting stents, 1182
- Uric acid, serum levels in patients with suspected or definite coronary artery disease undergoing elective vascular surgery, association between perioperative and late cardiovascular outcome and, 797
- VALsartan In Acute myocardial iNfarcTion trial (VALIANT) Registry
 - relation of mortality and failure to prescribe beta blockers acutely in patients with sustained ventricular tachycardia and ventricular fibrillation after acute myocardial infarction, 1427
- Valves, mechanical prosthetic
 - bileaflet aortic, Doppler-catheter discrepancies in patients with, 1383
 - perioperative bridging therapy with unfractionated heparin or low-molecular-weight heparin in patients on long-term oral anticoagulants with (from the REGIMEN Registry), 879
- Vascular surgery, elective
 - in patients with suspected or definite coronary artery disease and association between serum uric acid levels and perioperative and late cardiovascular outcome for, 797
 - revascularization in patients with multivessel coronary artery disease usefulness before, 809
- Vasovagal syncope, vasodilation in, water ingestion and, 1060
- Vena cava thrombosis, incidence in the United States, 927
- Ventricular arrhythmias
 - life-threatening, sleep-disordered breathing in heart failure patients with implantable cardioverter-defibrillator and, 1064
 - magnitude and prognosis in patients hospitalized with acute coronary syndromes associated with (from the GRACE Registry), 1577
- Ventricular dyssynchrony. *See* Interventricular dyssynchrony; Left ventricular dyssynchrony
- Ventricular fibrillation
 - as complication of acute myocardial infarction, thirty-year trends (1975–2005) in the magnitude, patient characteristics, and hospital outcomes of patients with, 1595
 - sustained, after acute myocardial infarction, relation of mortality and failure to prescribe beta blockers acutely in patients with (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry), 1427
- Ventricular pacing
 - managed, in pediatric patients and patients with congenital heart disease, 866
 - overdrive, diagnosis of accessory pathway using the His-His (H-H) and atrial-atrial (A-A) intervals of the first entrained cycle during, 197
- Ventricular septal defect, exercise capacity and participation of children with, 1079
- Ventricular tachycardia, sustained, after acute myocardial infarction, relation of mortality and failure to prescribe beta blockers acutely in patients with (from the VALsartan In Acute myocardial iNfarcTion trial [VALIANT] Registry), 1427
- Virtual histology, classification of intravascular ultrasound appearance of newly placed drug-eluting stents, 1182
- Waist circumference, body mass index in diagnosing metabolic syndrome and identifying apparently healthy subjects at increased risk of cardiovascular disease versus, 40
- Waist-to-hip ratio, impact on clinical outcomes in patients with ST-segment elevation acute myocardial infarction, body mass index and (from the Korean Acute Myocardial Infarction Registry), 957
- Warfarin, diagnostic safety of coronary angiography during uninterrupted therapeutic treatment with, 386
- Water, ingestion of, vasodilation in vasovagal syncope and, 1060
- Weightlifters, trained male, left ventricular diastolic function at rest and during isometric exercise in, 97
- Whites, comparison of cardiovascular risk factors for high brachial pulse pressure in blacks versus (Charleston Heart Study, Evans County Study, NHANES I and II Studies), 1514
- Women
 - asymptomatic, with systematic lupus erythematosus or rheumatoid arthritis, coronary artery calcium and C-reactive protein in, 755
 - cardiovascular risk with non-specific chest pain in (from the Women's Health Initiative Hormone Trials), 693
 - comparison of clinical features of arrhythmogenic right ventricular cardiomyopathy in men versus, 1252
 - with coronary artery disease, peripheral arterial response compared to mental stress in men versus, 970
 - defining normal distributions of coronary artery calcium in men and (from the Framingham Heart Study), 1136
 - hypertensive and normotensive postmenopausal, effect of soy nuts on adhesion molecules and inflammation markers in, 84
 - post cardiac catheterization contrast-induced acute kidney injury, age and, 1482
 - premenopausal, diabetes mellitus, hypothalamic hypoestrogenemia, and coronary artery disease in (from the National Heart, Lung, and Blood Institute Sponsored WISE Study), 150
 - with sedentary occupations and without known cardiovascular disease, endothelial function relation to cardiovascular risk in, 348
- Women's Health Initiative Hormone Trials, cardiovascular risk in women with non-specific chest pain, 693
- Worcester Heart Attack Study
 - thirty-year trends (1975–2005) in the magnitude, patient characteristics, and hospital outcomes of patients with acute myocardial infarction complicated by ventricular fibrillation, 1595
 - trends in prehospital delay in patients with acute myocardial infarction, 1589